

On Strong Convergence Theorems for END Sequences

Yamei Lu^{1,2}, Yan Shen^{1,*}, Rong Xu¹ and Xiaoqin Li¹

¹ School of Mathematical Science, Anhui University, Hefei 230039, P.R.China,

² Department of Mathematics, East China Normal University, Shanghai 200062, P.R.China.

Received February 16, 2016; Accepted March 6, 2016

Abstract. In this paper, we study the strong law of large numbers for a sequence of END random variables. Our results extend the corresponding ones for independent random variables and negatively orthant dependent (NOD, in short) random variables.

AMS subject classifications: 360F15

Key words: Convergence theorem, three series theorems, END random variable.

1 Introduction

Definition 1.1. ([1]) Random variables X_1, X_2, \dots are said to be extended negatively dependent (END, in short), if there exists a constant $M > 0$ such that both

$$P(X_1 \leq x_1, \dots, X_n \leq x_n) \leq M \prod_{i=1}^n P(X_i \leq x_i), \quad (1.1)$$

$$P(X_1 > x_1, \dots, X_n > x_n) \leq M \prod_{i=1}^n P(X_i > x_i) \quad (1.2)$$

hold for each $n \geq 2$ and all real numbers x_1, \dots, x_n .

The concept of END sequences was introduced by Liu [1]. When $M=1$, END random variables are negatively orthant dependent (NOD, in short) random variables, which was introduced by Joag-Dev and Proschan [2]. Some results for NOD sequences can be found in Ko and Kim [3], Fakoor and Azarnoosh [4], Ko *et al.* [5], Wu [6], Kim [7], Wu and Zhu [8]. As is mentioned in Liu [1], the END structure is substantially more comprehensive than the NOD structure in that it can reflect not only a negative dependence structure

*Corresponding author. *Email addresses:* 15155974615@163.com (Y. Lu), shenyan@ahu.edu.cn (Y. Shen), 530147340@qq.com (R. Xu), lixiaoqin1983@163.com (X. Li)

but also a positive one, to some extent. Liu [1] pointed out that the END random variables can be taken as negatively or positively dependent and provided some interesting examples to support this idea. Liu [1] obtained the precise large deviation results for END sequences. Liu [9] studied sufficient and necessary conditions for moderate deviations. Wu and Guan [10] discussed convergence properties of the partial sums for sequences of END random variables. Qiu *et al.* [11] obtained complete convergence for arrays of rowwise END random variables. For more details about strong convergence results for dependent sequence, one can refer to Sung [12], Wang *et al.* [13], Yang and Hu [14], Zhou *et al.* [15] and Zhou [16], and so forth.

The rest of the paper is organized as follows. In Section 2, some preliminary lemmas are presented. In Section 3, main results and their proofs are provided. Throughout the paper, let $I(A)$ be the indicator function of the set A . C denotes a positive constant not depending on n .

2 Preliminaries

The following lemmas will be needed in this paper.

Lemma 2.1 ([1]). *Let $\{X_n, n \geq 1\}$ be a sequence of END random variables, let f_1, f_2, \dots be all nondecreasing (or all nonincreasing) functions, then $\{f_n(X_n), n \geq 1\}$ is still a sequence of END random variables.*

Lemma 2.2 ([17]). *Let $\{X_n, n \geq 1\}$ be a sequence of END random variables. Assume that*

$$\sum_{n=1}^{\infty} \log^2 n \text{Var} X_n < \infty, \tag{2.1}$$

then $\sum_{n=1}^{\infty} (X_n - EX_n)$ converges almost surely.

By Lemmas 2.1 and 2.2, we can get the following three series theorems for END sequences. The proof is standard, so we omit it.

Lemma 2.3. *Let $\{X_n, n \geq 1\}$ be a sequence of END random variables. For some $c > 0$, denote $X_n^{(c)} = -cI(X_n < -c) + XI(|X_n| \leq c) + cI(X_n > c)$. If the following three conditions are satisfied:*

$$\sum_{n=1}^{\infty} P(|X_n| > c) < \infty, \tag{2.2}$$

$$\sum_{n=1}^{\infty} EX_n^{(c)} \text{ converges}, \tag{2.3}$$

$$\sum_{n=1}^{\infty} \log^2 n \text{Var} X_n^{(c)} < \infty, \tag{2.4}$$

then $\sum_{n=1}^{\infty} X_n$ converges almost surely.