The Neumann Problem of Complex Special Lagrangian Equations with Supercritical Phase

Chuanqiang Chen 1 , Xinan Ma 2,* and Wei Wei 2

¹ *Department of Applied Mathematics, Zhejiang University of Technology, Hangzhou, 310023, Zhejiang, China* ² *School of Mathematical Sciences, University of Science and Technology of China, Hefei, 230026, Anhui, China*

Received 22 August 2017; Accepted (in revised version) 8 June 2018

Abstract. Inspired by the Neumann problem of real special Lagrangian equations with supercritical phase, we consider the Neumann problem of complex special Lagrangian equations with supercritical phase in this paper, and establish the global *C* 2 estimates and the existence theorem by the method of continuity.

Key Words: Special Lagrangian equation, Neumann problem, supercritical phase.

AMS Subject Classifications: 35J60, 35B45

1 Introduction

As we all know, the real special Lagrangian equation is

$$
\arctan D^2 u = \Theta(x),\tag{1.1}
$$

where

$$
\arctan D^2 u := \arctan \lambda_1 + \arctan \lambda_2 + \dots + \arctan \lambda_n,
$$

 $\lambda = (\lambda_1, \lambda_2, \cdots, \lambda_n)$ are the eigenvalues of the Hessian matrix

$$
D^2 u = \left\{ \frac{\partial^2 u}{\partial x_i \partial x_j} \right\}_{1 \le i,j \le n'}
$$

and Θ is called the phase. In particular, $\Theta = \frac{(n-2)\pi}{2}$ $\frac{(-2)\pi}{2}$ is the critical phase, and if $\frac{(n-2)\pi}{2}$ < $\Theta(x) < \frac{n\pi}{2}$, Eq. (1.1) is called special Lagrangian equations with supercritical phase.

http://www.global-sci.org/ata/ 144 ©2019 Global-Science Press

[∗]Corresponding author. *Email addresses:* chuanqiangchen@zjut.edu.cn (C. Q. Chen), xinan@ustc.edu.cn (X. N. Ma), weisx001@mail.ustc.edu.cn (W. Wei)

The special Lagrangian equation (1.1) was introduced by Harvey-Lawson [13] in the study of calibrated geometries. Here Θ is a constant called the phase angle. In this case the graph $x \mapsto (x, Du(x))$ defines a calibrated, minimal submanifold of \mathbb{R}^{2n} . Since the work of Harvey-Lawson, special Lagrangian manifolds have gained wide interest, due in large part to their fundamental role in the Strominger-Yau-Zaslow description of mirror symmetry [26].

For the special Lagrangian equations with supercritical phase, Yuan obtained the interior C¹ estimate with Warren in [29] and the interior C² estimate with Wang in [28]. Recently Collins-Picard-Wu [8] obtained the existence theorem of the Dirichlet problem.

Moreover, for the Dirichlet problem of elliptic equations in \mathbb{R}^n , many results are known. For example, the Dirichlet problem of Laplace equation is studied in [7, 11], Caffarelli-Nirenberg-Spruck [2] and Ivochkina [15] solved the Dirichlet problem of Monge-Ampère equation, and Caffarelli-Nirenberg-Spruck [4] solved the Dirichlet problem of *k*-Hessian equation. After the pioneering works of Caffarelli et al., the Dirichlet problem of the general Hessian quotient equation was solved by Trudinger in [27]. For more information about the related subjects, we refer to the citations of [2, 4, 15].

Also, the Neumann or oblique derivative problem of partial differential equations was widely studied. For a priori estimates and the existence theorem of Laplace equation with Neumann boundary condition, we refer to the book [11]. Also, we can see the recent book written by Lieberman [21] for the Neumann and the oblique derivative problems of linear and quasilinear elliptic equations. In 1986, Lions-Trudinger-Urbas solved the Neumann problem of Monge-Ampère equation in the celebrated paper [23]. Recently, Ma-Qiu [24] solved the the Neumann problem of *k*-Hessian equations, and Chen-Zhang [6] generalized the result to the the Neumann problem of Hessian quotient equations. For the Neumann problem of special Lagrangian equations with supercritical phase, Chen-Ma-Wei [5] got the existence theorem. In [16, 17], Jiang-Trudinger studied the general oblique boundary value problems for augmented Hessian equations with some regular condition and some concavity condition.

At the same time, the complex equations have attracted a variety of mathematicians and many excellent works have been done. The complex Monge-Ampère equations are definitely one of the most important equations in partial differential equation and the geometry. In [1], Bedford and Taylor studied the Dirichlet problem of complex Monge-Ampère equations by using the Perron-Bremermann family method, and got the existence and uniqueness of the weak solutions and a global Lipschitz regularity for plurisubharmonic solution when Ω is a bounded strictly pseudoconvex domain in \mathbb{C}^n . In [3], Caffarelli et al. studied the classical solution on strongly pseudoconvex domains. Their work was extended to arbitrary bounded domains in **C***ⁿ* by Guan in [12] under some subsolution condition. Recently, Fu-Yau equation on compact Kähler manifolds arises much attention, which was introduced in [10]. The Fu-Yau equation was solved in dimension 2 by Fu-Yau and was recently extended to higher dimensions in some cases in [25] by Phong, Picard and Zhang. In high dimensions, the equation is actually a 2-Hessian type equation. Complex Hessian equations have been studied extensively by many authors

in recent years. On \mathbb{C}^n , the Dirichlet problem of complex Hessian equation was studied by Li [20] under some strict subsolution condition. On compact manifolds there were the existence of admissible solution in [9, 14]. For the Neumann problem of complex Monge-Ampère, Li proved the existence under some other conditions in [19].

Naturally, we want to know how about the problem of complex special Lagrangian equations of the form

$$
\begin{cases}\n\arctan\partial\overline{\partial}u = \Theta(z) & \text{in } \Omega, \\
D_{\nu}u = -\varepsilon u + \varphi(z) & \text{on } \partial\Omega,\n\end{cases}
$$
\n(1.2)

where Ω is a strictly pseudo-convex domain in \mathbb{C}^n , $\varepsilon > 0$ is constant, ν is the outer unit normal vector of *∂*Ω, *u* is a *C* ² and real valued function in Ω, Θ is a real valued function in Ω , $\partial \bar{\partial} u$ $=$ $\{\frac{\partial^2 u}{\partial z_i \partial \bar{z}}$ *∂zi∂z^j* }1≤*i*,*j*≤*n*, and arctan*∂* ¯*∂u* :=: arctan*λ*1+arctan*λ*2+···+arctan*λn*, where *λ*= $(\lambda_1, \lambda_2, \dots, \lambda_n)$ are the eigenvalues of the complex Hessian matrix $\partial \overline{\partial} u$. In the following, we denote

$$
z = (z_1, \dots, z_n) \in \overline{\Omega}, \quad z_j = t_j + \sqrt{-1}t_{n+j}, \quad t = (t_1, \dots, t_n, t_{n+1}, \dots, t_{2n}),
$$

\n
$$
\partial_j u = \frac{\partial u}{\partial z_j} = u_{z_j}, \quad \partial_j u = \frac{\partial u}{\partial z_j} = u_{z_j}, \quad \partial u = (\partial_1 u, \dots, \partial_n u),
$$

\n
$$
D_k u = \frac{\partial u}{\partial t_k}, \quad Du = (D_1 u, \dots, D_{2n} u),
$$

where $\sqrt{-1}$ is the imaginary unit. It is easy to know

$$
\partial_j u = \frac{1}{2} [D_j u - \sqrt{-1} D_{n+j} u], \quad |\partial u|^2 = \sum_{j=1}^n \partial_j u \overline{\partial_j u} = \frac{1}{4} |Du|^2.
$$

$$
\partial_{j\overline{j}} u = \frac{1}{4} [D_{jj} u + D_{n+j} u].
$$

In this paper, we establish global *C* 2 estimate of the Neumann problem of special Lagrangian equations with supercritical phase and obtain the existence theorem as follows:

Theorem 1.1. *Suppose* $\Omega \subset \mathbb{C}^n$ *is a* C^4 *strictly pseudoconvex domain and* $\varphi \in C^3(\partial \Omega)$ *.* Let $\Theta(z)\in C^2(\overline{\Omega})$ with $\frac{(n-2)\pi}{2}<\Theta(z)<\frac{n\pi}{2}$ in $\overline{\Omega}$ *. Moreover, if the domain* Ω satisfies a geometric *condition*

$$
2\kappa_{\min} + \varepsilon > 0,\tag{1.3}
$$

*where κ*min *is the smallest principal curvature of ∂*Ω*, then there exists a unique solution u* ∈ $C^{3,\alpha}(\overline{\Omega})$ of the Neumann problem of special Lagrangian equation (1.2).

As Li mentioned in his paper [19], because the Neumann boundary condition is not invariant under the holomorphic changes of variables, the strictly pesudoconvex domain may not be enough and we need the extra curvature condition of the domain.

Furthermore, we can obtain the existence theorem of the classical Neumann problem of complex special Lagrangian equation as below.

Theorem 1.2. *Suppose* Ω⊂**C***ⁿ is a C*⁴ *strictly convex domain and ν is outer unit normal vector* ∂f $\partial \Omega$ *. Let* $\varphi \in C^3(\partial \Omega)$ and $\Theta(z) \in C^2(\overline{\Omega})$ with $\frac{(n-2)\pi}{2} < \Theta(z) < \frac{n\pi}{2}$ in $\overline{\Omega}$ *. Then there exists a unique constant β such that the Neumann problem of special Lagrangian equation*

$$
\begin{cases}\n\arctan\partial\bar{\partial}u = \Theta(z) & \text{in } \Omega, \\
D_{\nu}u = \beta + \varphi(z) & \text{on } \partial\Omega,\n\end{cases}
$$
\n(1.4)

has admissible solutions $u\!\in\! C^{3,\alpha}(\overline{\Omega})$ *, which are unique up to a constant.*

The rest of the paper is organized as follows. In Section 2, we collect some properties of the special Lagrangian equation and establish the C^0 estimate for the Neumann problem of special Lagrangian equation. The *C* ¹ and *C* 2 estimates are established in Section 3, Section 4, respectively. At last, we prove Theorem 1.1 and Theorem 1.2 in Section 5.

2 Some properties and *C* ⁰ **estimate**

In this section, we give some properties of the special Lagrangian equation with supercritical phase and establish the C^0 estimate.

2.1 Some properties

Property 2.1. Suppose $\Omega \subset \mathbb{C}^n$ is a domain and $\Theta(z) \in C^0(\overline{\Omega})$ with $\frac{(n-2)\pi}{2} < \Theta(z) < \frac{n\pi}{2}$ in $\overline{\Omega}.$ Let $u\!\in\!\mathcal{C}^2(\Omega)$ be a solution of special Lagrangian equation

$$
\arctan \partial \bar{\partial} u = \Theta(z) \quad \text{in } \Omega. \tag{2.1}
$$

We assume $\lambda = (\lambda_1, \lambda_2, \cdots, \lambda_n)$ are the eigenvalues of the Hessian matrix $\partial \bar{\partial} u$ with $\lambda_1 \geq$ $\lambda_2 \cdots \geq \lambda_n$, then we have some properties:

$$
\lambda_1 \ge \dots \ge \lambda_{n-1} > 0,\tag{2.2a}
$$

$$
|\lambda_n| \le \lambda_{n-1},\tag{2.2b}
$$

$$
\sum_{i=1}^{n} \lambda_i > 0,\tag{2.2c}
$$

$$
|\lambda_n| < C_0,\tag{2.2d}
$$

.

where

$$
C_0 = \max \left\{ \tan \left(\frac{(n-1)\pi}{2} - \min_{\overline{\Omega}} \Theta \right), \tan \left(\frac{1}{n} \max_{\overline{\Omega}} \Theta \right) \right\}
$$

Proof. For any $i = 1, 2, \cdots, n$, we know arctan $\lambda_i \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Then we can get

$$
\arctan \lambda_{n-1} + \arctan \lambda_n = \Theta - \arctan \lambda_1 - \arctan \lambda_2 \cdots - \arctan \lambda_{n-2}
$$

$$
\geq \Theta - \frac{(n-2)\pi}{2} > 0.
$$

So $\lambda_{n-1} + \lambda_n > 0$, which implies (2.2a), (2.2b) and (2.2c) hold. Moreover,

$$
\arctan \lambda_n = \Theta - \arctan \lambda_1 - \arctan \lambda_2 - \dots - \arctan \lambda_{n-1} > \min_{\overline{\Omega}} \Theta - \frac{(n-1)\pi}{2}
$$

and

$$
\arctan \lambda_n < \frac{\arctan \lambda_1 + \arctan \lambda_2 + \dots + \arctan \lambda_n}{n} \le \frac{1}{n} \max_{\overline{\Omega}} \Theta,
$$

so we can get

$$
|\lambda_n| < \max\left\{\tan\left(\frac{(n-1)\pi}{2} - \min_{\overline{\Omega}} \Theta\right), \tan\left(\frac{1}{n} \max_{\overline{\Omega}} \Theta\right)\right\}.
$$

Thus, we complete the proof.

Property 2.2. Suppose $\Omega \subset \mathbb{C}^n$ is a domain and $\Theta(z) \in C^2(\overline{\Omega})$ with $\frac{(n-2)\pi}{2} < \Theta(z) < \frac{n\pi}{2}$ in $\overline{\Omega}.$ Let $u\in C^{4}(\Omega)$ be a solution of complex special Lagrangian equation (2.1). Then for any *ξ* ∈**S** 2*n*−1 , we have

$$
\sum_{ij=1}^{n} F^{i\bar{j}} D_{\xi\xi} \partial_{i\bar{j}} u \ge D_{\xi\xi} \Theta - A |D_{\xi}\Theta|^2 \quad \text{in} \quad \Omega,
$$
\n(2.3)

 \Box

where

$$
F^{i\bar{j}} = \frac{\partial \arctan \partial \bar{\partial} u}{\partial u_{i\bar{j}}} \quad \text{and} \quad A = \frac{2}{\tan\left(\min_{\overline{\Omega}} \Theta - \frac{(n-2)\pi}{2}\right)}.
$$

Proof. For any $z \in \Omega$, we can assume $\partial \overline{\partial} u$ is diagonal with $\lambda_i = u_{i\overline{i}}$, since (2.3) is invariant under rotating the coordinates. Then we have

$$
F^{i\bar{j}} =: \frac{\partial \arctan \partial \bar{\partial} u}{\partial u_{i\bar{j}}} = \begin{cases} \frac{1}{1 + \lambda_i^2}, & \text{if } i = j, \\ 0, & \text{if } i \neq j, \end{cases}
$$

and

$$
F^{i\bar{j},k\bar{l}} =: \frac{\partial^2 \arctan \partial \bar{\partial} u}{\partial u_{i\bar{j}} \partial u_{k\bar{l}}} = \begin{cases} -\frac{2\lambda_i}{(1+\lambda_i^2)^2}, & \text{if } i=j=k=l, \\ -\frac{\lambda_i + \lambda_j}{(1+\lambda_i^2)(1+\lambda_j^2)}, & \text{if } i=l, j=k, i \neq j, \\ 0, & \text{otherwise.} \end{cases}
$$

From the complex special Lagrangian equation (2.1), we know

$$
\sum_{ij=1}^n F^{i\bar{j}} D_{\xi} \partial_{i\bar{j}} u = D_{\xi} \Theta
$$

and

$$
\sum_{ij=1}^{n} F^{i\bar{j}} D_{\xi\xi} \partial_{i\bar{j}} u = D_{\xi\xi} \Theta - \sum_{ijkl=1}^{n} F^{i\bar{j},k\bar{l}} D_{\xi} \partial_{i\bar{j}} u D_{\xi} \partial_{k\bar{l}} u
$$

\n
$$
= D_{\xi\xi} \Theta - \sum_{i=1}^{n} F^{i\bar{i},l\bar{i}} (D_{\xi} \partial_{i\bar{i}} u)^{2} - \sum_{i \neq j} F^{i\bar{j},l\bar{i}} |D_{\xi} \partial_{i\bar{j}} u|^{2}
$$

\n
$$
\geq D_{\xi\xi} \Theta - \sum_{i=1}^{n} F^{i\bar{i},l\bar{i}} (D_{\xi} \partial_{i\bar{i}} u)^{2}.
$$
 (2.4)

From the concavity lemma (Lemma 2.2 in [8]), we know

$$
-\sum_{i=1}^{n} F^{i\overline{i}, i\overline{i}} (D_{\xi} \partial_{i\overline{i}} u)^{2} \geq -\frac{2}{\tan\left(\min_{\overline{\Omega}} \Theta - \frac{(n-2)\pi}{2}\right)} \left(\sum_{i=1}^{n} F^{i\overline{i}} D_{\xi} \partial_{i\overline{i}} u\right)^{2}
$$

$$
= -\frac{2}{\tan\left(\min_{\overline{\Omega}} \Theta - \frac{(n-2)\pi}{2}\right)} |D_{\xi} \Theta|^{2}.
$$
(2.5)

Hence (2.3) holds.

2.2 *C* ⁰ **estimate**

The C⁰ estimate is easy. For completeness*,* we produce a proof here following the idea of Lions-Trudinger-Urbas [23] and Ma-Qiu [24].

 $\bf{Theorem 2.3.}$ $Suppose \ \Omega \subset \mathbb{C}^n$ is a C^1 bounded domain and $\varphi \! \in \! C^0(\partial \Omega)$. Let $\Theta(z) \! \in \! C^0(\overline{\Omega})$ with $\frac{(n-2)\pi}{2}$ $<$ $\Theta(z)$ $<$ $\frac{n\pi}{2}$ *in* $\overline{\Omega}$ *and* u \in $C^2(\Omega)$ \cap $C^1(\overline{\Omega})$ *be the solution of complex special Lagrangian equation* (1.2)*, then we have*

$$
-M_0 \le u(z) \le \frac{1}{\varepsilon} \max_{\partial \Omega} |\varphi| \quad \text{in} \quad \Omega,\tag{2.6}
$$

where

$$
M_0 = \frac{1}{\varepsilon} \max_{\partial \Omega} |\varphi| + \frac{2B}{\varepsilon} \operatorname{diam}(\Omega) + \operatorname{Bdiam}(\Omega)^2 \quad \text{and} \quad B = \tan\left(\frac{1}{n} \max_{\overline{\Omega}} \Theta\right) < +\infty.
$$

Proof. From (2.2c), we know *u* is subharmonic. So the maximum of *u* is attained at a boundary point *z*0∈*∂*Ω. Then we can get

$$
0 \le D_{\nu}u(z_0) = -\varepsilon u(z_0) + \varphi(z_0). \tag{2.7}
$$

 \Box

Hence

$$
\max_{\overline{\Omega}} u = u(z_0) \le \frac{\varphi(z_0)}{\varepsilon} \le \frac{1}{\varepsilon} \max_{\partial \Omega} |\varphi|.
$$
 (2.8)

Without loss of generality, we assume $0 \in \Omega$ and denote

$$
B=\tan\left(\frac{1}{n}\max_{\overline{\Omega}}\Theta\right)<+\infty.
$$

Then we have

$$
\arctan\partial\bar{\partial}u = \Theta \le \max_{\overline{\Omega}} \Theta = \arctan\partial\bar{\partial}(B|z|^2). \tag{2.9}
$$

By the comparison principle, we know *u*−*B*|*z*| ² attains its minimum at a boundary point *z*1∈*∂*Ω. Then

$$
0 \ge D_{\nu}(u - B|z|^2)|_{z=z_1} = D_{\nu}u(z_1) - BD_{\nu}(|z|^2)|_{z=z_1}
$$

= $-\varepsilon u(z_1) + \varphi(z_1) - 2B \cdot t \cdot \nu|_{z=z_1}$
 $\ge -\varepsilon u(z_1) - \max_{\partial \Omega} |\varphi| - 2B \text{diam}(\Omega).$ (2.10)

Hence

$$
\min_{\overline{\Omega}} u \ge \min_{\overline{\Omega}} (u - B|z|^2) = u(z_1) - B|z_1|^2
$$

$$
\ge -\frac{1}{\varepsilon} \max_{\partial \Omega} |\varphi| - \frac{2B}{\varepsilon} \text{diam}(\Omega) - B \text{diam}(\Omega)^2.
$$
 (2.11)

Thus, we complete the proof.

3 Global gradient estimate

In this section, we prove the global gradient estimate by following the idea of Li [19].

Theorem 3.1. *Suppose* $\Omega \subset \mathbb{C}^n$ *is a* C^3 *bounded domain satisfying* (1.3) *and* $\varphi \in C^2(\partial \Omega)$ *.* Let $\Theta(z) \in C^1(\overline{\Omega})$ *with* $\frac{(n-2)\pi}{2} < \Theta(z) < \frac{n\pi}{2}$ in $\overline{\Omega}$ and $u \in C^3(\Omega) \cap C^2(\overline{\Omega})$ be a solution of complex *special Lagrangian equation* (1.2)*, then we have*

$$
\sup_{\overline{\Omega}}|Du| \le M_1,\tag{3.1}
$$

 \Box

where M_1 *depends on n,* Ω *,* $\kappa_{\min} + \varepsilon$ *,* $|u|_{C^0}$ *,* $\max_{\overline{O}} \Theta$ *,* $\min_{\overline{O}} \Theta$ *and* $|\varphi|_{C^2}$ *.*

Proof. In order to prove (3.1), it suffices to prove

$$
D_{\xi}u(z) \le M_1, \quad \forall (z,\xi) \in \overline{\Omega} \times \mathbb{S}^{2n-1}.
$$
 (3.2)

For any $(z,\xi)\in\overline{\Omega}\times\mathbb{S}^{2n-1}$, denote

$$
W(z,\xi) = D_{\xi}u(z) - \langle v,\xi \rangle(-\varepsilon u + \varphi(z)) + K_0 u^2 + K_1 |z|^2,
$$
\n(3.3)

where K_0 and K_1 are large constants to be determined later, and ν is a $C^2(\overline{\Omega})$ extension of the outer unit normal vector field on *∂*Ω. We still use *ϕ*(*z*) to denote a *C* 2 extension of *ϕ* from *∂*Ω to Ω. Assume *W* achieves its maximum at (*z*0,*ξ*0)∈Ω×**S** 2*n*−1 . It is easy to know $D_{\xi_0}u(z_0) > 0$. We claim *z*₀ ∈ ∂Ω. Otherwise, if *z*₀ ∈ Ω, we rotate the coordinates such that $\{\partial_{i\bar{j}}u(z_0)\}\$ is diagonal with $\lambda_i = u_{i\bar{i}}$ and $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n$. Then we have

$$
F^{i\bar{j}} =: \frac{\partial \arctan \partial \bar{\partial} u}{\partial u_{i\bar{j}}} = \begin{cases} \frac{1}{1 + \lambda_i^2}, & \text{if } i = j, \\ 0, & \text{if } i \neq j. \end{cases}
$$

Hence we can get from Property 2.1

$$
F^{1\bar{1}} \le F^{2\bar{2}} \le \dots \le F^{n\bar{n}},
$$

\n
$$
F^{n\bar{n}} = \frac{1}{1 + \lambda_n^2} \ge c_0 > 0,
$$

\n
$$
F^{i\bar{i}} u_{i\bar{i}} = \frac{\lambda_i}{1 + \lambda_i^2} \in \left(-\frac{1}{2}, \frac{1}{2}\right),
$$

where

$$
c_0 = \frac{1}{1 + \max\{\tan\left(\frac{(n-1)\pi}{2} - \min\Theta\right), \tan\left(\frac{1}{n}\max\Theta\right)\}^2}.
$$

For fixed $\xi = \xi_0$, $W(z, \xi_0)$ achieves its maximum at the same point $z_0 \in \Omega$ and we can easily get that at z_0 ,

$$
0 \geq F^{i\bar{j}} \partial_{i\bar{j}} W = F^{i\bar{i}} \left[\partial_{i\bar{i}} D_{\xi_0} u - \langle v, \xi_0 \rangle_{i\bar{i}} (-\varepsilon u + \varphi) - \langle v, \xi_0 \rangle (-\varepsilon u_{i\bar{i}} + \varphi_{i\bar{i}}) - \langle v, \xi_0 \rangle_i (-\varepsilon u_{\bar{i}} + \varphi_{\bar{i}}) - \langle v, \xi_0 \rangle_{\bar{i}} (-\varepsilon u_i + \varphi_i) + 2K_0 u_i u_{\bar{i}} + 2K_0 u u_{i\bar{i}} + K_1 \right] \n\geq D_{\xi_0} \Theta + (2K_0 - C_1) \sum_{i=1}^n F^{i\bar{i}} u_i u_{\bar{i}} \n+ [2K_0 u + \varepsilon \langle v, \xi_0 \rangle] \sum_{i=1}^n F^{i\bar{i}} u_{i\bar{i}} + (K_1 - C_1) \sum_{i=1}^n F^{i\bar{i}},
$$
\n(3.4)

where C_1 is a positive constant depending only on *n*, ε , $|u|_{C^0}$, $|\varphi|_{C^2}$ and $|v|_{C^2}$.

Take $K_0 = \frac{C_1}{2}$ and $K_1 = 2C_1 + \frac{2|D\Theta| + n[2K_0|u| + \varepsilon]}{c_0}$ $\frac{\Gamma[2N_0|u|+\epsilon]}{c_0}$. Using *n* ∑ $F^{i\bar{i}}u_{i\bar{i}} \in \left(-\frac{n}{2}\right)$ $\frac{n}{2}, \frac{n}{2}$ 2 , *n* ∑ $F^{i\bar{i}} > F^{n\bar{n}} \ge c_0$,

i=1

we can get

$$
0 \ge F^{i\bar{j}} \partial_{i\bar{j}} W
$$

\n
$$
\ge -|D_{\xi_0} \Theta| - \frac{n}{2} [2K_0|u| + \varepsilon] + (2K_0 - C_1) \sum_{i=1}^n F^{i\bar{i}} u_i u_{\bar{i}} + (K_1 - C_1) \sum_{i=1}^n F^{i\bar{i}}
$$

\n
$$
> 0.
$$
\n(3.5)

i=1

This is a contradiction. So $z_0 \in \partial \Omega$. Then we continue our proof in the following three cases.

(a) If ξ_0 is normal at $z_0 \in \partial \Omega$, then

$$
W(z_0, \xi_0) = K_0 u^2 + K_1 |z_0|^2 \le C_2.
$$

Then we can easily get (3.2).

(b) If ξ_0 is non-tangential at $z_0 \in \partial \Omega$, then we can write $\xi_0 = \alpha \tau + \beta \nu$, where $\tau \in \mathbb{S}^{2n-1}$ is tangential at *z*₀, that is $\langle \tau, \nu \rangle = 0$, $\alpha = \langle \xi_0, \tau \rangle > 0$, $\beta = \langle \xi_0, \nu \rangle < 1$, and $\alpha^2 + \beta^2 = 1$. Then we have

$$
W(z_0, \xi_0) = \beta D_{\tau} u + K_0 u^2 + K_1 |z_0|^2
$$

\n
$$
\leq \beta W(z_0, \xi_0) + (1 - \beta)(K_0 u^2 + K_1 |z_0|^2),
$$
\n(3.6)

so

$$
W(z_0,\xi_0) \le K_0 u^2 + K_1 |z_0|^2 \le C_3.
$$

Then we can easily get (3.2).

(c) If *ξ*⁰ is tangential at *z*0∈*∂*Ω, we may assume that the outer normal direction of Ω at *z*⁰ is $(0, \dots, 0, 1)$. By a rotation, we assume that $\xi_0 = (1, \dots, 0) = e_1$. Then we have

$$
0 \le D_{\nu}W(z_0,\xi_0) = D_{\nu}D_1u - D_{\nu}\langle \nu,\xi_0 \rangle(-\varepsilon u + \varphi) + 2K_0u \cdot D_{\nu}u + K_1D_{\nu}|z|^2
$$

\n
$$
\le D_{\nu}D_1u + C_4
$$

\n
$$
= D_1D_{\nu}u - D_1\nu_kD_ku + C_4.
$$
\n(3.7)

By the boundary condition, we know

$$
D_1 D_v u = D_1(-\varepsilon u + \varphi) \le -\varepsilon W(z_0, \xi_0) + C_5. \tag{3.8}
$$

Following the argument of [19], we can get

$$
-D_1 \nu_k D_k u \le -\kappa_{\min} W(z_0, \xi_0) + C_6. \tag{3.9}
$$

From (1.3), it holds $\kappa_{\min}+\varepsilon > 0$. So

$$
W(z_0, \xi_0) \le \frac{C_4 + C_5 + C_6}{\kappa_{\min} + \varepsilon}.
$$
\n(3.10)

Then we can easily get (3.2).

Remark 3.1. To prove Theorem 1.2, we need to consider the complex special Lagrangian equation (1.2) with *ε*→0. Under the assumptions of Theorem 1.2, that is, Ω ⊂**C***ⁿ* is a *C* 4 strictly convex domain, $\varphi \in C^3(\partial \Omega)$, $\Theta(z) \in C^2(\overline{\Omega})$ and $\frac{(n-2)\pi}{2} < \Theta(z) < \frac{n\pi}{2}$ in $\overline{\Omega}$, and for any sufficiently small *ε*, we can obtain the following proof of Theorem 2.3

$$
\sup_{\overline{\Omega}} \varepsilon |u^{\varepsilon}| \leq \widetilde{M}_0,\tag{3.11}
$$

for the solution u^ε of (1.2), where M_0 depends on *n, diam*(Ω), max Θ , and $|\varphi|_{C^0}.$ Moreover, we can choose test function

$$
W(z,\xi) = D_{\xi}u^{\varepsilon}(z) - \langle v,\xi \rangle (-\varepsilon u^{\varepsilon} + \varphi(z)) + K_0(\varepsilon u^{\varepsilon})^2 + K_1|z|^2,
$$

and following the proof of Theorem 3.1, we can choose *K*⁰ and *K*¹ independent of *ε*, and obtain the global gradient estimate

$$
\sup_{\overline{\Omega}}|Du^{\varepsilon}|\leq \widetilde{M}_1,\tag{3.12}
$$

where M_1 depends on *n*, Ω , $\max_{\overline{O}} \Theta$, $\min_{\overline{O}} \Theta$, $|\Theta|_{C^1}$ and $|\varphi|_{C^2}$, and is independent of *ε*. It is Ω Ω easy to know

$$
\sup_{\overline{\Omega}} \left| u^{\varepsilon} - \frac{1}{|\Omega|} \int_{\Omega} u^{\varepsilon} \right| \le \widetilde{M}_1. \tag{3.13}
$$

4 Global second derivatives estimate

We now come to the a priori estimate of global second derivatives. Firstly, we prove the estimate of double normal second derivatives on boundary, and then we complete the proof of global second derivatives estimate.

 \Box

4.1 Estimate of double normal second derivatives on boundary

Theorem 4.1. *Suppose* $\Omega \subset \mathbb{C}^n$ *is a* C^4 *strictly pseudoconvex domain and* $\varphi \in C^3(\partial \Omega)$ *.* Let $\Theta(z)\!\in\! C^2(\overline{\Omega})$ with $\frac{(n-2)\pi}{2}\!<\!\Theta(z)\!<\!\frac{n\pi}{2}$ in $\overline{\Omega}$ and $u\!\in\! C^4(\Omega)\!\cap\! C^3(\overline{\Omega})$ be the solution of complex *special Lagrangian equation* (1.2)*, then we have*

$$
\max_{\partial \Omega} |D_{\nu\nu} u| \le \overline{M_2},\tag{4.1}
$$

 w here $\overline{M_2}$ depends on n , Ω , $\max_{\overline{\Omega}} \Theta$, $|u|_{C^1}$, $\min_{\overline{\Omega}} \Theta$, $|\Theta|_{C^2}$ and $|\varphi|_{C^3}$.

Proof. Since Ω is a C^4 strictly pseudoconvex domain, there is a strictly plurisubharmonic defining function r ∈ $C^4(\overline{\Omega})$ such that

$$
|Dr| = 1 \qquad \text{on } \partial \Omega,
$$
 (4.2a)

$$
\partial \bar{\partial} r \ge k_0 I_n \quad \text{in } \overline{\Omega}, \tag{4.2b}
$$

where k_0 is a positive constant depending only on Ω and I_n is the $n \times n$ identity matrix.

Let $z_0 \in \partial \Omega$ be an arbitrary point. By a shift and a rotation of the coordinates ${z_1, \dots, z_n}$, we can assume that $z_0 = 0$, $\partial_{z_i} r(0) = 0$ for $i < n$, and $D_{t_n} r(0) = −1$, $D_{t_{2n}} r(0) = 0$. In $\overline{B}(0,\delta)\bigcap\Omega$, a sufficiently small neighborhood of z_0 , we can get by the Taylor expansion of *r* up to second order

$$
r(z) = -Re\left(z_n - \sum_{ij=1}^n a_{ij} z_i z_j\right) + \sum_{ij=1}^n b_{ij} z_i \overline{z_j} + O(|z|^3),\tag{4.3}
$$

where $\{b_{i\bar{j}}\} = \partial \bar{\partial} r(0)$ is positive definite. We now introduce new coordinates $z' = \psi(z)$ of the form

$$
z'_i = z_i
$$
 for $i < n$, $z'_n = z_n - \sum_{i,j=1}^n a_{ij} z_i z_j$. (4.4)

In $\psi(\overline{B}(0,\delta)\bigcap \overline{\Omega})$, we have

$$
r(z)|_{z=\psi^{-1}(z')} = -Re z'_n + \sum_{ij=1}^n b_{i\bar{j}} z'_i \overline{z'_j} + O(|z'|^3), \qquad (4.5)
$$

Denote

$$
r_0(z') = -Rez'_n + \sum_{ij=1}^n b_{ij}z'_i \overline{z'_j},
$$

\n
$$
B_j(z') = \Big(\sum_{i=1}^n [a_{ij} + a_{ji}]z_i\Big)\Big|_{z=\psi^{-1}(z')} , \quad j = 1, \dots, n,
$$

\n
$$
A_j(z') = B_j(1 - B_n)^{-1} \frac{\partial r_0(z')}{\partial \overline{z'_n}}, \quad j = 1, \dots, n-1.
$$

It is easy to know $|B_j|$ $=$ $\mathcal{O}(|z'|)$ for j $=$ $1,$ \cdots , n , and A_j is holomorphic in z' \in $\psi(\overline{B}(0,\delta)\cap \overline{\Omega}).$ Following the calculations in [19], we know the Neumann boundary condition in z' coordinates

$$
4Re\left(\langle\partial_{z'}u,\partial_{z'}r_0\rangle-\sum_{j=1}^{n-1}A_j\partial_{z'_j}u\right)=\phi(z',u)+\mathcal{O}(|z'|^3),\tag{4.6}
$$

where

$$
\phi(z', u) = |1 - B_n(z')|^{-2} (-\varepsilon u + \varphi(z))|_{z = \psi^{-1}(z')}.
$$

Following the idea of [19], we choose the auxiliary function

$$
h(z') = 4Re\left[\langle \partial_{z'} u, \partial_{z'} r_0 \rangle - \sum_{j=1}^{n-1} A_j \partial_{z'_j} u\right] - \phi(z', u) + Kr(z)|_{z=\psi^{-1}(z')} - K_1 Re(z'_n),\tag{4.7}
$$

where $K_1 > 0$ is sufficiently large so that

$$
h < 0 \quad \text{on } \psi(\partial(B(0,\delta)) - \partial \Omega),
$$

and

$$
h = -K_1 Re(z'_n) + \mathcal{O}(|z'|^2) \le 0 \quad \text{on } \psi(\partial(B(0,\delta)) \cap \partial \Omega).
$$

Let

$$
F^{i\bar{j}} = \frac{\partial \arctan \partial_z \overline{\partial_z} u}{\partial u_{z_i \overline{z_j}}}, \quad G^{i\bar{j}} = \frac{\partial \arctan \partial_z \overline{\partial_z} u}{\partial u_{z'_i \overline{z'_j}}}.
$$

It is easy to know

$$
G^{i\bar{j}} = F^{p\bar{q}} \left(\frac{\partial z_i'}{\partial z_p} \right) \overline{\left(\frac{\partial z_j'}{\partial z_q} \right)}.
$$

For any $z' \in \psi(B(0,\delta) \cap \Omega)$, we can get

$$
G^{i\bar{j}}\partial_{z_{i}^{i}\overline{z_{j}^{\prime}}}h=2G^{i\bar{j}}\partial_{z_{i}^{i}\overline{z_{j}^{\prime}}}(\partial_{z_{k}^{i}}u\partial_{\overline{z_{k}^{\prime}}}r_{0}+\partial_{\overline{z_{k}^{\prime}}}u\partial_{z_{k}^{\prime}}r_{0})-2G^{i\bar{j}}\partial_{z_{i}^{i}\overline{z_{j}^{\prime}}}\left(\sum_{k=1}^{n-1}A_{k}\partial_{z_{k}^{i}}u+\overline{A_{k}}\partial_{\overline{z_{k}^{\prime}}}u\right)\\-G^{i\bar{j}}\partial_{z_{i}^{i}\overline{z_{j}^{\prime}}}\phi+KG^{i\bar{j}}\partial_{z_{i}^{i}\overline{z_{j}^{\prime}}}r_{0}\\=2G^{i\bar{j}}(\partial_{z_{k}^{i}}u_{z_{i}^{i}\overline{z_{j}^{\prime}}}\partial_{\overline{z_{k}^{\prime}}}r_{0}+\partial_{\overline{z_{k}^{\prime}}}u_{z_{i}^{i}\overline{z_{j}^{\prime}}}\partial_{z_{k}^{i}}r_{0}+u_{z_{k}^{i}\overline{z_{j}^{\prime}}}\partial_{z_{i}^{i}\overline{z_{k}^{\prime}}}r_{0}+u_{\overline{z_{k}^{\prime}}}u_{\overline{z_{k}^{\prime}}}\partial_{z_{k}^{i}\overline{z_{j}^{\prime}}}r_{0})\\-2\sum_{k=1}^{n-1}G^{i\bar{j}}(A_{k}\partial_{z_{k}^{i}}u_{z_{i}^{i}\overline{z_{j}^{\prime}}}+\overline{A_{k}}\partial_{\overline{z_{k}^{\prime}}}u_{z_{i}^{i}\overline{z_{j}^{\prime}}}+\partial_{z_{i}^{\prime}}A_{k}u_{z_{k}^{i}\overline{z_{j}^{\prime}}}+\overline{\partial_{z_{j}^{\prime}}A_{k}u_{z_{i}^{i}\overline{z_{k}^{\prime}}}})\\-G^{i\bar{j}}[|1-B_{n}(z^{\prime})|^{-2}(-\varepsilon u_{z_{i}^{i}\overline{z_{j}^{\prime}}}+\varphi_{z_{i}^{i}\overline{z_{j}^{\prime}}})+\partial_{z_{i}^{i}\overline{z_{j}^{\prime}}}(|1-B_{n}(z^{\prime})|^{-2
$$

156 C. Q. Chen, X. N. Ma and W. Wei / Anal. Theory Appl., **35** (2019), pp. 144-162

$$
=2\left[\partial_{z'_{k}}\Theta\partial_{\overline{z'_{k}}}r_{0}+\partial_{\overline{z'_{k}}}\Theta\partial_{z'_{k}}r_{0}\right]+2F^{p\bar{q}}\left(u_{z'_{k}\overline{z_{q}}}\partial_{z_{p}\overline{z'_{k}}}r_{0}+u_{\overline{z'_{k}}z_{p}}\partial_{z'_{k}\overline{z_{q}}}r_{0}\right) -2\sum_{k=1}^{n-1}\left[A_{k}\partial_{z'_{k}}\Theta+\overline{A_{k}}\partial_{\overline{z'_{k}}}\Theta+F^{p\bar{q}}\left(\partial_{z_{p}}A_{k}u_{z'_{k}\overline{z_{q}}}+\overline{\partial_{z_{q}}A_{k}}u_{z_{p}\overline{z'_{k}}}\right)\right] -F^{p\bar{q}}\left[|1-B_{n}(z')|^{-2}\left(-\varepsilon u_{z_{p}\overline{z_{q}}}+\varphi_{z_{p}\overline{z_{q}}}\right)+\partial_{z_{p}\overline{z_{q}}}\left(|1-B_{n}(z')|^{-2}\right)\left(-\varepsilon u+\varphi\right)\right] +KF^{p\bar{q}}\partial_{z_{p}\overline{z_{q}}}r \geq Kk_{0}\sum_{p=1}^{n}F^{p\bar{p}}-C_{7},
$$
\n(4.8)

where $F^{p\bar{q}}u_{z_k'\overline{z_q}} = F^{p\bar{q}}u_{z_m\overline{z_q}}\frac{\partial z_m}{\partial z_k'}$ $\frac{\partial z_m}{\partial z'_k}$ and $F^{p\bar{q}}u_{z_p\overline{z'_k}} = F^{p\bar{q}}u_{z_p\overline{z_m}}(\frac{\partial z_m}{\partial z'_k})$ $\frac{d z_m}{\partial z'_k}$) are bounded by rotating the coordinates $\{z_1, \dots, z_n\}$ such that $\partial_z \overline{\partial_z} u$ is diagonal. From (2.2d), we know

$$
\sum_{p=1}^{n} F^{p\bar{p}} \ge c_0,
$$
\n(4.9)

where

$$
c_0 = \frac{1}{1 + \max\{\tan\left(\frac{(n-1)\pi}{2} - \min_{\overline{\Omega}} \Theta\right), \tan\left(\frac{1}{n} \max_{\overline{\Omega}} \Theta\right)\}^2}.
$$

Taking $K = \frac{C_7}{k_0c}$ $\frac{C_7}{k_0c_0}$, we can have $G^{i\bar{j}}\partial_{z_i^{\prime}\overline{z_j^{\prime}}}h≥0$ in $\psi(B(0,\delta)\cap\Omega)$. By the maximum principle, we know $h(z')$ achieves its maximum at $z' = 0$. Hence $h(z')|_{z' = \psi(z)}$ achieves its maximum at $z_0 = 0$. Thus

$$
0 \le D_{\nu}h(0) \le D_{\nu\nu}u(z_0) + C_8. \tag{4.10}
$$

So we have $D_{\nu\nu}u(z_0) \geq -C_8$.

The same argument for

$$
h_1(z') = 4Re\left[\langle \partial_{z'} u, \partial_{z'} r_0 \rangle - \sum_{j=1}^{n-1} A_j \partial_{z'_j} u\right] - \phi(z', u) - Kr(z)|_{z=\psi^{-1}(z')} + K_1 Re(z'_n)
$$
(4.11)

can give

$$
D_{\nu\nu}u(z_0)\leq C_9.\t\t(4.12)
$$

This completes the estimates of the double normal derivative on the boundary. \Box

4.2 Estimate of global second derivatives

Theorem 4.2. Suppose $\Omega \subset \mathbb{C}^n$ is a C^4 strictly pseudoconvex domain satisfying (1.3) and $\varphi \in$ $C^3(\partial\Omega)$ *. Let* $\Theta(z) \in C^2(\overline{\Omega})$ *with* $\frac{(n-2)\pi}{2} < \Theta(z) < \frac{n\pi}{2}$ in $\overline{\Omega}$ and $u \in C^4(\Omega) \cap C^3(\overline{\Omega})$ be a solution

of special Lagrangian equation (1.2)*, then we have*

$$
\sup_{\overline{\Omega}}|D^2u|\leq M_2,\tag{4.13}
$$

 w *here* M_2 *depends on n,* Ω *,* $2\kappa_{min} + \varepsilon$ *,* $\max_{\overline{\Omega}} \Theta$ *,* $\min_{\overline{\Omega}} \Theta$ *,* $|u|_{C^1}$ *,* $|\Theta|_{C^2}$ *and* $|\varphi|_{C^3}$ *.*

Proof. From (2.2c), we know

$$
\Delta u = 4 \sum_{i=1}^{n} u_{i\bar{i}} = 4(\lambda_1 + \dots + \lambda_n) > 0.
$$

By the argument in Li's [19], we know that we only need to prove that

$$
D_{\zeta\zeta}u(z) \le M_2, \quad \forall (z,\zeta) \in \bar{\Omega} \times \mathbb{S}^{2n-1}.
$$
 (4.14)

As the real case in [23], we use the auxiliary function

$$
Q(z,\zeta) = D_{\zeta\zeta}u - v(z,\zeta) + K_1|z|^2 + |Du|^2,
$$
\n(4.15)

where $v(z,\zeta) = 2\langle \zeta,\nu \rangle \langle \zeta', D\varphi - \varepsilon Du - D_1 u D \nu^1 \rangle = a_1 D_1 u + b$, $v = (v^1, v^2, \dots, v^{2n}) \in \mathbb{S}^{2n-1}$ is a *C* 3 (Ω) extension of the outer unit normal vector field on *∂*Ω

$$
\zeta' = \zeta - \langle \zeta, \nu \rangle \nu, \quad a_l = -2 \langle \zeta, \nu \rangle \langle \zeta', D\nu^l \rangle - 2\varepsilon \langle \zeta, \nu \rangle (\zeta')^l, \quad b = 2 \langle \zeta, \nu \rangle \langle \zeta', D\varphi \rangle,
$$

and $K_1 > 0$ is to be determined later.

For any $z \in \Omega$, we rotate the coordinates such that $\{\partial_{i\bar{j}}u(z)\}$ is diagonal with $\lambda_i = u_{i\bar{i}}$ and $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n$. Then we have

$$
F^{i\bar{j}} =: \frac{\partial \arctan \partial \bar{\partial} u}{\partial u_{i\bar{j}}} = \begin{cases} \frac{1}{1 + \lambda_i^2}, & \text{if } i = j, \\ 0, & \text{if } i \neq j. \end{cases}
$$

Hence we can get from Property 2.1

$$
F^{1\bar{1}} \le F^{2\bar{2}} \le \dots \le F^{n\bar{n}},
$$

\n
$$
F^{n\bar{n}} = \frac{1}{1 + \lambda_n^2} \ge c_0 > 0,
$$

\n
$$
F^{i\bar{i}} u_{i\bar{i}} = \frac{\lambda_i}{1 + \lambda_i^2} \in \left(-\frac{1}{2}, \frac{1}{2}\right),
$$

where $c_0 = \frac{1}{1 + \max\left\{\tan\left(\frac{(n-1)\pi}{2} - \min_{\overline{\Omega}} \Theta\right), \tan\left(\frac{1}{n} \max_{\overline{\Omega}} \Theta\right)\right\}}^2$.

For any fixed $\zeta \in \mathbb{S}^{n-1}$, we have

$$
F^{i\bar{j}}\partial_{i\bar{j}}Q(z,\zeta) = F^{i\bar{j}}\partial_{i\bar{j}}D_{\zeta\zeta}u - F^{i\bar{i}}[\partial_{i\bar{i}}a_{k}D_{k}u + \partial_{\bar{i}}a_{k}\partial_{i}D_{k}u + \partial_{i}a_{k}\partial_{\bar{i}}D_{k}u + a_{k}\partial_{i\bar{i}}D_{k}u + \partial_{i\bar{i}}b]
$$

+2F<sup>i\bar{i}\partial_{i}D_{k}u\partial_{\bar{i}}D_{k}u + 2D_{k}\Theta D_{k}u + K_{1}\sum_{i=1}^{n}F^{i\bar{i}}

$$
\geq D_{\zeta\zeta}\Theta - A|D_{\zeta}\Theta|^{2} - (|D^{2}a_{k}||Du| + |D^{2}b|)\sum_{i=1}^{n}F^{i\bar{i}} - F^{i\bar{i}}(\partial_{\bar{i}}a_{k}\partial_{i}D_{k}u + \partial_{i}a_{k}\partial_{\bar{i}}D_{k}u)
$$

$$
-|a_{k}D_{k}\Theta| + 2F^{i\bar{i}}\partial_{i}D_{k}u\partial_{\bar{i}}D_{k}u - 2|D_{k}\Theta D_{k}u| + K_{1}\sum_{i=1}^{n}F^{i\bar{i}}.
$$
(4.16)</sup>

It is easy to know

$$
\sum_{i=1}^{n} F^{i\bar{i}} > F^{n\bar{n}} \ge c_0,
$$
\n
$$
2F^{i\bar{i}}\partial_i D_k u \partial_{\bar{i}} D_k u - F^{i\bar{i}} (\partial_{\bar{i}} a_k \partial_i D_k u + \partial_i a_k \partial_{\bar{i}} D_k u) \ge -F^{i\bar{i}} \partial_i a_k \partial_{\bar{i}} a_k.
$$

Choose

$$
K_1 = 2[|D^2 a_k||Du| + |D^2 b|] + \frac{2[|D^2 \Theta| + A|D\Theta|^2 + |a_k||D\Theta| + 2|D\Theta||Du|]}{c_0},
$$

then we can get

$$
F^{i\bar{j}}\partial_{i\bar{j}}Q(z,\zeta) \ge 0. \tag{4.17}
$$

So maxQ(*z*,*ζ*) attains its maximum on ∂Ω. Hence $\frac{max}{\overline{\Omega} \times S^{2n-1}}$ Q(*z*,*ζ*) attains its maximum at some point $z_0 \in \partial \Omega$ and some direction $\zeta_0 \in \mathbb{S}^{2n-1}$.

Then we continue our proof in the following two cases following the idea of [19].

(a) If ζ_0 is non-tangential at $z_0 \in \partial \Omega$, then we can write $\zeta_0 = \alpha \tau + \beta \nu$, where $\tau \in \mathbb{S}^{2n-1}$ is tangential at z_0 , that is $\langle \tau,\nu\rangle$ = 0, α = $\langle \zeta_0,\tau\rangle$, β = $\langle \zeta_0,\nu\rangle$ \neq 0, and $\alpha^2+\beta^2$ = 1. Then we have

$$
D_{\zeta_0\zeta_0}u(z_0)
$$

= $\alpha^2 D_{\tau\tau}u(z_0) + \beta^2 D_{\nu\nu}u(z_0) + 2\alpha\beta D_{\tau\nu}u(z_0)$
= $\alpha^2 D_{\tau\tau}u(z_0) + \beta^2 D_{\nu\nu}u(z_0) + 2(\zeta_0 \cdot \nu)[\zeta_0 - (\zeta_0 \cdot \nu)\nu] \cdot [D\varphi - \varepsilon Du - D_l u D\nu^l],$ (4.18)

hence

$$
Q(z_0, \zeta_0) = \alpha^2 Q(z_0, \tau) + \beta^2 Q(z_0, \nu).
$$
\n(4.19)

From the definition of $Q(z_0, \zeta_0)$, we know

$$
Q(z_0, \zeta_0) \le Q(z_0, \nu) \le C_{10} \tag{4.20}
$$

and we can prove (4.14).

(b) If
$$
\xi_0
$$
 is tangential at $z_0 \in \partial\Omega$, then we have

$$
0 \le D_{\nu}Q(z_0,\zeta_0) = D_{\nu}D_{\zeta_0\zeta_0}u - D_{\nu}a_lD_lu - a_lD_{\nu}D_lu - D_{\nu}b + 2D_kuD_{\nu}D_ku + K_1D_{\nu}|z|^2
$$

\n
$$
\le D_{\nu}D_{\zeta_0\zeta_0}u + [2D_ku - a_k]D_{\nu}D_ku + C_{11}.
$$
\n(4.21)

By the boundary condition, we know

$$
D_{\nu}D_{\zeta_{0}\zeta_{0}}u = D_{\zeta_{0}\zeta_{0}}D_{\nu}u - (D_{\zeta_{0}\zeta_{0}}\nu^{k})D_{k}u - 2(D_{\zeta_{0}}\nu^{k})D_{\zeta_{0}}D_{k}u
$$

\n
$$
= D_{\zeta_{0}\zeta_{0}}(-\varepsilon u + \varphi) - (D_{\zeta_{0}\zeta_{0}}\nu^{k})D_{k}u - 2(D_{\zeta_{0}}\nu^{k})D_{\zeta_{0}}D_{k}u
$$

\n
$$
\leq -\varepsilon Q(z_{0},\zeta_{0}) + C_{12} - 2(D_{\zeta_{0}}\nu^{k})D_{\zeta_{0}}D_{k}u.
$$
\n(4.22)

Following the argument of [19], we can get

$$
|D_{\nu}D_{k}u| \leq C_{13},
$$

-2(D_{ζ0}v^k)D_{ζ0}D_ku $\leq -2\kappa_{\min}Q(z_0,\zeta_0)+C_{14}.$

From (1.3), it holds $2\kappa_{\min}+\varepsilon > 0$. So

$$
Q(z_0,\zeta_0) \le \frac{C_{11} + C_{12} + (2|Du| + |a_k|)C_{13} + C_{14}}{2\kappa_{\min} + \varepsilon}.
$$
\n(4.23)

Then we can easily get (4.14).

Remark 4.1. As the discussions in Remark 3.1, to prove Theorem 1.2, we need to consider the complex special Lagrangian equation (1.2) with $\varepsilon \rightarrow 0$. Under the assumptions of Theorem 1.2, and for any sufficiently small *ε*, we can obtain the following proof of Theorem 4.1 and Theorem 4.2,

$$
\sup_{\overline{\Omega}} |D^2 u^{\varepsilon}| = \sup_{\overline{\Omega}} \left| D^2 \left(u^{\varepsilon} - \frac{1}{|\Omega|} \int_{\Omega} u^{\varepsilon} \right) \right| \le \widetilde{M}_2,\tag{4.24}
$$

where M_2 depends on *n*, Ω, $\max_{\overline{\Omega}}$ Θ, $\min_{\overline{\Omega}}$ θ, $|\Theta|_{C^2}$ and $|\varphi|_{C^3}$, and is independent of *ε*.

5 Existence of the boundary problems

In this section we complete the proofs of the Theorem 1.1 and Theorem 1.2.

5.1 Proof of Theorem 1.1

For the Neumann problem of complex special Lagrangian equation (1.2), we have established the C^0 , C^1 and C^2 estimates in Section 2, Section 3, Section 4, respectively. By the global C^2 priori estimate*,* the special Lagrangian equation (1.2) is uniformly elliptic in $\overline{\Omega}.$

 \Box

From the concavity Lemma (Lemma 2.2 in [8]), we know −*e* −*A*arctan*∂* ¯*∂u* is concave with respect to *∂* ¯*∂u*, where *A* is defined in Property 2.2. Following the discussions in [22], we can get the global Hölder estimates of second derivative,

$$
|u|_{C^{2,\alpha}(\overline{\Omega})} \leq C,\tag{5.1}
$$

where *C* and *α* depend on *n*, Ω , $\max_{\overline{\Omega}} \Theta$, $\min_{\overline{\Omega}} \Theta$, $|\Theta|_{C^2}$ and $|\varphi|_{C^3}$. From (5.1), one also obtains $C^{3,\alpha}(\overline{\Omega})$ estimates by differentiating Eq. (1.2) and applies the Schauder theory for linear uniformly elliptic equations.

Applying the method of continuity (see [21]), the existence of the classical solution holds. By the standard regularity theory of uniformly elliptic partial differential equations, we can obtain the higher regularity.

5.2 Proof of Theorem 1.2

By a similar proof of Theorem 1.1, we know there exists a unique solution $u^{\varepsilon} \! \in \! C^{3,\alpha}(\overline{\Omega})$ to (1.2) for any small $\varepsilon > 0$. Let $v^{\varepsilon} = u^{\varepsilon} - \frac{1}{|\Omega|} \int_{\Omega} u^{\varepsilon}$, and it is easy to know that v^{ε} satisfies

$$
\begin{cases}\n\arctan\partial\bar{\partial}v^{\varepsilon} = \Theta(z) & \text{in } \Omega, \\
D_{\nu}(v^{\varepsilon}) = -\varepsilon v^{\varepsilon} - \frac{1}{|\Omega|} \int_{\Omega} \varepsilon u^{\varepsilon} + \varphi(z) & \text{on } \partial\Omega.\n\end{cases}
$$
\n(5.2)

By the global gradient estimate (3.12), it is easy to know $\epsilon \sup|Du^{\epsilon}|\to 0$. Hence there is a Ω $\text{constant } \beta$ and a function $v\!\in\! C^2(\overline{\Omega})$, such that

$$
-\varepsilon u^{\varepsilon} \to \beta, \quad -\varepsilon v^{\varepsilon} \to 0, \quad -\frac{1}{|\Omega|} \int_{\Omega} \varepsilon u^{\varepsilon} \to \beta, \quad v^{\varepsilon} \to v,
$$

uniformly in $C^2(\overline{\Omega})$ as $\varepsilon \!\to\! 0.$ It is easy to verify that v is a solution of

$$
\begin{cases}\n\arctan\partial\bar{\partial}v = \Theta(z) & \text{in } \Omega, \\
D_{\nu}v = \beta + \varphi(z) & \text{on } \partial\Omega.\n\end{cases}
$$
\n(5.3)

If there is another function v_1 \in $C^2(\overline{\Omega})$ and another constant β_1 such that

$$
\begin{cases}\n\arctan\partial\bar{\partial}v_1 = \Theta(z) & \text{in } \Omega, \\
D_v(v_1) = \beta_1 + \varphi(z) & \text{on } \partial\Omega.\n\end{cases}
$$
\n(5.4)

Applying the maximum principle and Hopf Lemma, we can know $\beta = \beta_1$ and $v - v_1$ is a constant. By the standard regularity theory of uniformly elliptic partial differential equations, we can obtain the higher regularity.

Acknowledgements

Research of the first author was supported by ZJNSF No. LY17A010022 and NSFC No. 11771396. Research of the second author and third author was supported by NSFC No. 11471188 and Wu Wen-Tsun Key Laboratory of Mathematics in USTC. Research of the third author was supported by China Scholarship Council.

References

- [1] E. Bedford and J. E. Fornaess, Counterexamples to regularity for the complex Monge-Ampere equation, Invent. Math., 50 (1979), 129–134. `
- [2] L. Caffarelli, L. Nirenberg and J. Spruck, The Dirichlet problem for nonlinear second order elliptic equations I, Monge-Ampère equations, Commun. Pure Appl. Math., 37 (1984), 369-402.
- [3] L. A. Caffarelli, J. J. Kohn, L. Nirenberg and J. Spruck, The Dirichlet problem for nonlinear second-order elliptic equations, II. Complex Monge-Ampère and uniformly elliptic equations, Commun. Pure Appl. Math., 38 (1985), 209–252.
- [4] L. Caffarelli, L. Nirenberg and J. Spruck. The Dirichlet problem for nonlinear second order elliptic equations III, Functions of the eigenvalues of the Hessian, Acta Math., 155 (1985), 261–301.
- [5] C. Q. Chen, X. N. Ma and W. Wei, The Neumann problem of special Lagrangian equations with supercritical phase, Preprint, 2017.
- [6] C. Q. Chen and D. K. Zhang, The Neumann problem for Hessian quotient equations, Preprint, 2016.
- [7] Y. Z. Chen and L. C. Wu, Second Order Elliptic Equations and Elliptic Systems, Am. Math. Soc., Providence, RI, 1998.
- [8] T. C. Collins, S. Picard and X. Wu, Concavity of the Lagrangian phase operator and applications, Calc. Var., 56 (2017), 89.
- [9] S. Dinew and S. Kolodziej, Liouville and Calabi-Yau type theorems for complex Hessian equations, Am. J. Math., 139(2) (2017), 403–415.
- [10] J. X. Fu and S. T. Yau, The theory of superstring with flux on non-Kähler manifolds and the complex Monge-Ampère equation, J. Diff. Geom., 78(3) (2008), 369-428.
- [11] D. Gilbarg and N. Trudinger, Elliptic Partial Differential Equations of Second Order, Grundlehren der Mathematischen Wissenschaften, Vol. 224. Springer-Verlag, Berlin-New York, 1977. x+401 pp. ISBN: 3-540-08007-4.
- [12] B. Guan, The Dirichlet problem for complex Monge-Ampère equations and regularity of the pluri-complex Green function, Comm. Anal. Geom., 6 (1998), 687–703; a correction, 8 (2000), 213–218.
- [13] R. Harvey and B. Lawson, Calibrated geometries, Acta Math., 148(1) (1982), 47–157.
- [14] Z. L. Hou, X. N. Ma and D. M. Wu, A second order estimate for complex Hessian equations on a compact Kähler manifold, Math. Res. Lett., 17 (2010), 547-561.
- [15] N. Ivochkina, Solutions of the Dirichlet problem for certain equations of Monge-Ampere ` type (in Russian), Mat. Sb., 128 (1985), 403–415: English translation in Math. USSR Sb., 56 (1987).
- [16] F. Jiang and N. S. Trudinger, Oblique boundary value problems for augmented Hessian equations I, Bull. Math. Sci., 8 (2018), 353–411.
- [17] F. Jiang and N. S. Trudinger, Oblique boundary value problems for augmented Hessian equations II, Nonlinear Anal. Theory Methods Appl., 154 (2017), 148–173.
- [18] S. Kolodziej and N.-C. Nguyen, Weak solutions of complex Hessian equations on compact Hermitian manifolds, Compos. Math., 152 (2016), 2221–2248.
- [19] S. Y. Li, On the Neumann problems for Complex Monge-Ampere equations, Indiana Univ. ` Math. J., 43 (1994), 1099–1122.
- [20] S. Y. Li, On the Dirichlet problems for symmetric function equations of the eigenvalues of the complex Hessian, Asian J. Math., 8 (2004), 087–106.
- [21] G. Lieberman, Oblique Boundary Value Problems for Elliptic Equations, World Scientific Publishing, (2013).
- [22] G. Lieberman and N. Trudinger, Nonlinear oblique boundary value problems for nonlinear elliptic equations, Trans. Amer. Math. Soc., 295 (1986), 509-546.
- [23] P. L. Lions, N. Trudinger and J. Urbas, The Neumann problem for equations of Monge-Ampere type, Commun. Pure Appl. Math., 39 (1986), 539–563. `
- [24] X. N. Ma and G. H. Qiu, The Neumann Problem for Hessian Equations, Commun. Math. Phys., 366 (2019), 1–28.
- [25] D. H. Phong, S. Picard and X. W. Zhang, The Fu-Yau equation with negative slope parameter, Invent. Math., 209 (2017), 541–576.
- [26] A. Stromginer, S.-T. Yau and E. Zaslow, Mirror symmetry is *T*-duality, Nuclear Phys. B, 479 (1996), 243–259.
- [27] N. S. Trudinger, On the Dirichlet problem for Hessian equations, Acta Math., 175 (1995), 151–164.
- [28] D. K. Wang and Y. Yuan, Hessian estimates for special Lagrangian equations with critical and supercritical phases in general dimensions, Amer. J. Math., 136 (2014), 481–499.
- [29] M. Warren and Y. Yuan, Hessian and gradient estimates for three dimensional special Lagrangian equations with large phase, Amer. J. Math., 132 (2010), 751–770.