

Application of Homotopy Perturbation Transform Method for Solving Linear and Nonlinear Klein-Gordon Equations

Jagdev Singh¹, Devendra Kumar² and Sushila Rathore³

¹Department of Mathematics, Jagan Nath University, Village-Rampura, Tehsil-Chaksu, Jaipur-303901, Rajasthan, India

²Department of Mathematics, Jagan Nath Gupta Institute of Engineering and Technology, Jaipur-302022, Rajasthan, India

³Department of Physics, Jagan Nath University, Village-Rampura, Tehsil-Chaksu, Jaipur-303901, Rajasthan, India

(Received November 7, 2011, accepted December 2, 2011)

Abstract. In this paper, the homotopy perturbation transform method (HPTM) has been applied to obtain the solution of the linear and nonlinear Klein-Gordon equations. The homotopy perturbation transform method is a combined form of the Laplace transform method with the homotopy perturbation method. This scheme finds the solution without any discretization or restrictive assumptions and avoids the round-off errors. The fact that this technique solves nonlinear problems without using Adomian's polynomials can be considered as a clear advantage of this technique over the decomposition method. The results reveal that the proposed algorithm is very efficient, simple and can be applied to other nonlinear problems.

Mathematics Subject Classification: 35C05, 35C10.

Key words and Phrases: Laplace transform method, Homotopy perturbation method, Linear and Nonlinear Klein-Gordon equations, He's Polynomials.

1. Introduction

Nonlinear phenomena have important effects on applied mathematics, physics and related to engineering; many such physical phenomena are modeled in terms of nonlinear partial differential equations. For example, the Klein-Gordon equations which are of the form

$$u_{tt}(x, t) - u_{xx}(x, t) + au(x, t) = h(x, t), \quad (1)$$

with initial conditions

$$u(x, 0) = f(x), \quad u_t(x, 0) = g(x), \quad (2)$$

appears in quantum field theory, relativistic physics, dispersive wave-phenomena, plasma physics, nonlinear optics and applied physical sciences. Several techniques including finite difference, collocation, finite element, inverse scattering, decomposition and variational iteration using Adomian's polynomials have been used to handle such equations [5, 6, 7, 48]. He [1-4] developed the homotopy perturbation technique for solving such physical problems. In recent years, many research workers have paid attention to study the solutions of nonlinear partial differential equations by using various methods. Among these are the Adomian decomposition method (ADM) [8], He's semi-inverse method [9], the tanh method, the homotopy perturbation method (HPM), the differential transform method and the variational iteration method (VIM) [10-17]. He [25-38] developed the homotopy perturbation method (HPM) by merging the standard homotopy and perturbation for solving various physical problems. The Laplace transform is totally incapable of handling nonlinear equations because of the difficulties that are caused by the nonlinear terms. Various ways have been proposed recently to deal with these nonlinearities such as the Adomian decomposition method

¹ E-mail address: jagdevsinghrathore@gmail.com;

² E-mail address: devendra.maths@gmail.com;

³ E-mail address: sushila.jag@gmail.com.

[39] and the Laplace decomposition algorithm [40-44]. Furthermore, the homotopy perturbation method is also combined with the well-known Laplace transformation method [45] and the variational iteration method [46] to produce a highly effective technique for handling many nonlinear problems. In a recent paper Khan and Wu [23] proposed the homotopy perturbation transform method (HPTM) for solving the nonlinear equations. It is worth mentioning that this method is an elegant combination of the Laplace transformation, the homotopy perturbation method and He's polynomials and is mainly due to Ghorbani [20, 21]. The homotopy perturbation transform method provides the solution in a rapid convergent series which may lead to the solution in a closed form. The advantage of this method is its capability of combining two powerful methods for obtaining exact solutions for nonlinear equations. In this article, we apply the homotopy perturbation transform method (HPTM) for solving the linear and nonlinear Klein-Gordon equations to show the simplicity and straight forwardness of the method.

2. Homotopy perturbation transform method (HPTM)

To illustrate the basic idea of this method, we consider a general nonlinear partial differential equation with the initial conditions of the form:

$$D u(x, t) + R u(x, t) + N u(x, t) = g(x, t), \quad (3)$$

$$u(x, 0) = h(x), \quad u_t(x, 0) = f(x),$$

where D is the second order linear differential operator $D = \partial^2 / \partial t^2$, R is the linear differential operator of less order than D , N represents the general nonlinear differential operator and $g(x, t)$ is the source term. Taking the Laplace transform (denoted in this paper by L) on both sides of eq. (3), we get

$$L[D u(x, t)] + L[R u(x, t)] + L[N u(x, t)] = L[g(x, t)]. \quad (4)$$

Using the differentiation property of the Laplace transform, we have

$$L[u(x, t)] = \frac{h(x)}{s} + \frac{f(x)}{s^2} + \frac{1}{s^2} L[g(x, t)] - \frac{1}{s^2} L[R u(x, t)] - \frac{1}{s^2} L[N u(x, t)]. \quad (5)$$

Operating with the Laplace inverse on both sides of eq. (5) gives

$$u(x, t) = G(x, t) - L^{-1} \left[\frac{1}{s^2} L[R u(x, t) + N u(x, t)] \right], \quad (6)$$

where $G(x, t)$ represents the term arising from the source term and the prescribed initial conditions. Now we apply the homotopy perturbation method

$$u(x, t) = \sum_{n=0}^{\infty} p^n u_n(x, t) \quad (7)$$

and the nonlinear term can be decomposed as

$$N u(x, t) = \sum_{n=0}^{\infty} p^n H_n(u), \quad (8)$$

for some He's polynomials $H_n(u)$ (see [21, 46]) that are given by

$$H_n(u_0, u_1, \dots, u_n) = \frac{1}{n!} \frac{\partial^n}{\partial p^n} \left[N \left(\sum_{i=0}^{\infty} p^i u_i \right) \right]_{p=0}, \quad n = 0, 1, 2, 3, \dots \quad (9)$$

Substituting eq. (7) and eq. (8) in eq. (6), we get

$$\sum_{n=0}^{\infty} p^n u_n(x, t) = G(x, t) - p \left(L^{-1} \left[\frac{1}{s^2} L \left[R \sum_{n=0}^{\infty} p^n u_n(x, t) + \sum_{n=0}^{\infty} p^n H_n(u) \right] \right] \right), \quad (10)$$

which is the coupling of the Laplace transform and the homotopy perturbation method using He's polynomials. Comparing the coefficient of like powers of p , the following approximations are obtained.

$$\begin{aligned}
 p^0 : u_0(x, t) &= G(x, t), \\
 p^1 : u_1(x, t) &= -L^{-1} \left[\frac{1}{s^2} L \left[R u_0(x, t) + H_0(u) \right] \right], \\
 p^2 : u_2(x, t) &= -L^{-1} \left[\frac{1}{s^2} L \left[R u_1(x, t) + H_1(u) \right] \right], \\
 p^3 : u_3(x, t) &= -L^{-1} \left[\frac{1}{s^2} L \left[R u_2(x, t) + H_2(u) \right] \right], \\
 &\vdots
 \end{aligned} \tag{11}$$

3. Numerical Applications

In this section, we use homotopy perturbation transform method (HPTM) in solving the linear and nonlinear Klein-Gordon equations.

Example 3.1. Consider the following linear Klein-Gordon equation

$$u_{tt}(x, t) - u_{xx}(x, t) + u(x, t) = 0, \tag{12}$$

with the initial conditions

$$u(x, 0) = 0, u_t(x, 0) = x. \tag{13}$$

Applying the Laplace transform on both sides of eq. (12) subject to the initial conditions (13), we have

$$L[u(x, t)] = \frac{x}{s^2} + \frac{1}{s^2} L[u_{xx}(x, t) - u(x, t)]. \tag{14}$$

The inverse of Laplace transform implies that

$$u(x, t) = xt + L^{-1} \left[\frac{1}{s^2} L[u_{xx}(x, t) - u(x, t)] \right]. \tag{15}$$

Now, applying the homotopy perturbation method, we get

$$\sum_{n=0}^{\infty} p^n u_n(x, t) = xt + p \left(L^{-1} \left[\frac{1}{s^2} L \left[\left(\sum_{n=0}^{\infty} p^n u_n(x, t) \right)_{xx} - \sum_{n=0}^{\infty} p^n u_n(x, t) \right] \right] \right). \tag{16}$$

Comparing the coefficients of like powers of p , we have

$$\begin{aligned}
 p^0 : u_0(x, t) &= xt, \\
 p^1 : u_1(x, t) &= L^{-1} \left[\frac{1}{s^2} L \left[(u_0)_{xx} - u_0 \right] \right] = \frac{-xt^3}{3!}, \\
 p^2 : u_2(x, t) &= -L^{-1} \left[\frac{1}{s^2} L \left[(u_1)_{xx} - u_1 \right] \right] = \frac{xt^5}{5!}, \\
 &\vdots
 \end{aligned} \tag{17}$$

Therefore the solution $u(x, t)$ is given by

$$u(x, t) = x \left(t - \frac{t^3}{3!} + \frac{t^5}{5!} - \dots \right), \quad (18)$$

in series form, and

$$u(x, t) = x \sin t, \quad (19)$$

in closed form.

Example 3.2. Consider the following linear Klein-Gordon equation

$$u_{tt}(x, t) - u_{xx}(x, t) + u(x, t) = 2 \sin x, \quad (20)$$

with the initial conditions

$$u(x, 0) = \sin x, \quad u_t(x, 0) = 1. \quad (21)$$

Applying the Laplace transform on both sides of eq. (20) subject to the initial conditions (21), we have

$$L[u(x, t)] = \frac{2 \sin x}{s^3} + \frac{\sin x}{s} + \frac{1}{s^2} + \frac{1}{s^2} L[u_{xx}(x, t) - u(x, t)]. \quad (22)$$

The inverse of Laplace transform implies that

$$u(x, t) = t^2 \sin x + \sin x + t + L^{-1} \left[\frac{1}{s^2} L[u_{xx}(x, t) - u(x, t)] \right]. \quad (23)$$

Now, applying the homotopy perturbation method, we get

$$\sum_{n=0}^{\infty} p^n u_n(x, t) = t^2 \sin x + \sin x + t + p \left(L^{-1} \left[\frac{1}{s^2} L \left[\left(\sum_{n=0}^{\infty} p^n u_n(x, t) \right)_{xx} - \sum_{n=0}^{\infty} p^n u_n(x, t) \right] \right] \right). \quad (24)$$

Comparing the coefficients of like powers of p , we have

$$p^0 : u_0(x, t) = t^2 \sin x + \sin x + t,$$

$$p^1 : u_1(x, t) = L^{-1} \left[\frac{1}{s^2} L[(u_0)_{xx} - u_0] \right] = -t^2 \sin x - \frac{t^3}{3!} - \frac{t^4}{3!} \sin x, \quad (25)$$

$$p^2 : u_2(x, t) = L^{-1} \left[\frac{1}{s^2} L[(u_1)_{xx} - u_1] \right] = \frac{t^4}{3!} \sin x + \frac{t^6}{90} \sin x + \frac{t^5}{5!},$$

⋮

Therefore the solution $u(x, t)$ is given by

$$u(x, t) = \sin x + \left(t - \frac{t^3}{3!} + \frac{t^5}{5!} - \dots \right) = \sin x + \sin t. \quad (26)$$

Example 3.3. Consider the following nonlinear Klein-Gordon equation

$$u_{tt}(x, t) - u_{xx}(x, t) + u^2(x, t) = x^2 t^2, \quad (27)$$

with the initial conditions

$$u(x,0) = 0, u_t(x,0) = x. \quad (28)$$

Applying the Laplace transform on both sides of eq. (27) subject to the initial conditions (28), we have

$$L[u(x,t)] = \frac{2x^2}{s^5} + \frac{x}{s^2} + \frac{1}{s^2} L[u_{xx}(x,t) - u^2(x,t)]. \quad (29)$$

The inverse of Laplace transform implies that

$$u(x,t) = \frac{x^2 t^4}{12} + xt + L^{-1} \left[\frac{1}{s^2} L[u_{xx}(x,t) - u^2(x,t)] \right]. \quad (30)$$

Now, applying the homotopy perturbation method, we get

$$\sum_{n=0}^{\infty} p^n u_n(x,t) = \frac{x^2 t^4}{12} + xt + p \left(L^{-1} \left[\frac{1}{s^2} L \left[\left(\sum_{n=0}^{\infty} p^n u_n(x,t) \right)_{xx} - \sum_{n=0}^{\infty} p^n H_n(u) \right] \right] \right), \quad (31)$$

where $H_n(u)$ are He's polynomial [21, 46] that represents the nonlinear terms. The first few components of He's polynomials, are given by

$$\begin{aligned} H_0(u) &= (u_0)^2, \\ H_1(u) &= 2u_0 u_1, \\ &\vdots \end{aligned} \quad (32)$$

Comparing the coefficients of like powers of p , we have

$$\begin{aligned} p^0 : u_0(x,t) &= \frac{x^2 t^4}{12} + xt, \\ p^1 : u_1(x,t) &= L^{-1} \left[\frac{1}{s^2} L \left[(u_0)_{xx} - H_0(u) \right] \right] \\ &= \frac{t^6}{180} - \frac{x^4 t^{10}}{12960} - \frac{x^3 t^7}{252} - \frac{x^2 t^4}{12}, \end{aligned} \quad (33)$$

$$\begin{aligned} p^2 : u_2(x,t) &= L^{-1} \left[\frac{1}{s^2} L \left[(u_1)_{xx} - H_1(u) \right] \right] \\ &= -\frac{x^2 t^{12}}{71280} - \frac{11xt^9}{22680} - \frac{t^6}{180} + \frac{x^6 t^{16}}{18662400} + \frac{11x^4 t^{10}}{45360} \\ &\quad + \frac{383x^5 t^{13}}{15921360} + \frac{x^3 t^7}{252}, \\ &\vdots \end{aligned}$$

Therefore the solution $u(x,t)$ is given by

$$u(x,t) = xt. \quad (34)$$

Example 3.4. Consider the following nonlinear Klein-Gordon equation

$$u_{tt}(x,t) - u_{xx}(x,t) + u^2(x,t) = 2x^2 - 2t^2 + x^4 t^4, \quad (35)$$

with the initial conditions

$$u(x,0) = 0, u_t(x,0) = 0. \quad (36)$$

Applying the Laplace transform on both sides of eq. (35) subject to the initial conditions (36), we have

$$L[u(x,t)] = \frac{2x^2}{s^3} - \frac{4}{s^5} + \frac{24x^4}{s^7} + \frac{1}{s^2} L[u_{xx}(x,t) - u^2(x,t)]. \quad (37)$$

The inverse of Laplace transform implies that

$$u(x,t) = x^2 t^2 - \frac{t^4}{6} + \frac{x^4 t^6}{30} + L^{-1} \left[\frac{1}{s^2} L[u_{xx}(x,t) - u^2(x,t)] \right]. \quad (38)$$

Now, applying the homotopy perturbation method, we get

$$\sum_{n=0}^{\infty} p^n u_n(x,t) = x^2 t^2 - \frac{t^4}{6} + \frac{x^4 t^6}{30} + p \left(L^{-1} \left[\frac{1}{s^2} L \left[\left(\sum_{n=0}^{\infty} p^n u_n(x,t) \right)_{xx} - \sum_{n=0}^{\infty} p^n H_n(u) \right] \right] \right), \quad (39)$$

where $H_n(u)$ are He's polynomial [21, 46] that represents the nonlinear terms.

In a similar manner as above the solution $u(x,t)$ is given by

$$u(x,t) = x^2 t^2. \quad (40)$$

Example 3.5. Consider the following nonlinear Klein-Gordon equation

$$u_{tt}(x,t) - u_{xx}(x,t) + u^2(x,t) = 6xt(x^2 - t^2) + x^6 t^6, \quad (41)$$

with the initial conditions

$$u(x,0) = 0, u_t(x,0) = 0. \quad (42)$$

Applying the Laplace transform on both sides of eq. (41) subject to the initial conditions (42), we have

$$L[u(x,t)] = \frac{6x^3}{s^4} - \frac{36x}{s^6} + x^6 \frac{8!}{s^{11}} + \frac{1}{s^2} L[u_{xx}(x,t) - u^2(x,t)]. \quad (43)$$

The inverse of Laplace transform implies that

$$u(x,t) = x^3 t^3 - \frac{3xt^5}{10} + \frac{x^6 t^{10}}{90} + L^{-1} \left[\frac{1}{s^2} L[u_{xx}(x,t) - u^2(x,t)] \right]. \quad (44)$$

Now, applying the homotopy perturbation method, we get

$$\sum_{n=0}^{\infty} p^n u_n(x,t) = x^3 t^3 - \frac{3xt^5}{10} + \frac{x^6 t^{10}}{90} + p \left(L^{-1} \left[\frac{1}{s^2} L \left[\left(\sum_{n=0}^{\infty} p^n u_n(x,t) \right)_{xx} - \sum_{n=0}^{\infty} p^n H_n(u) \right] \right] \right), \quad (45)$$

where $H_n(u)$ are He's polynomial [21, 46] that represents the nonlinear terms.

In a similar manner as above the solution $u(x,t)$ is given by

$$u(x, t) = x^3 t^3. \quad (46)$$

4. Conclusions

In this paper, the homotopy perturbation transform method (HPTM) has been successfully applied to find the solution of the linear and nonlinear Klein-Gordon equations with initial conditions. The method is reliable and easy to use. The results show that the homotopy perturbation transform method is powerful and efficient technique in finding exact and approximate solutions for nonlinear differential equations. It is worth mentioning that the method is capable of reducing the volume of the computational work as compared to the classical methods while still maintaining the high accuracy of the numerical result; the size reduction amounts to an improvement of the performance of the approach. The fact that the HPTM solves nonlinear problems without using Adomian's polynomials is a clear advantage of this technique over the decomposition method. In conclusion, the HPTM may be considered as a nice refinement in existing numerical techniques and might find the wide applications.

5. References

- [1] J.H. He. Homotopy perturbation technique. *Computer Methods in Applied Mechanics and Engineering*. 1999, **178**: 257–262.
- [2] N.H. Sweilam, M.M. Khader. Exact solutions of some coupled nonlinear partial differential equations using the homotopy perturbation method. *Computers & Mathematics with Applications*. 2009, **58**: 2134–2141.
- [3] J. Saberi-Nadjafi, A. Ghorbani. He's homotopy perturbation method: an effective tool for solving nonlinear integral and integro-differential equations. *Computers & Mathematics with Applications*. 2009, **58**: 1345–1351.
- [4] A. Yildirim. An Algorithm for Solving the Fractional Nonlinear Schrödinger Equation by Means of the Homotopy Perturbation Method. *International Journal of Nonlinear Science and Numerical Simulation*. 2009, **10**: 445-451.
- [5] S. Abbasbandy. Numerical solutions of nonlinear Klein-Gordon equation by variational iteration method. *Internat. J. Meth. Engrg*. 2007, **70**: 876-881.
- [6] S.T. Mohyud-Din, M.A. Noor, K.I. Noor. some relatively new techniques for nonlinear problems. *Math. Porb. Eng.* Article ID 234849, 25 pages, doi:10.1155/2009/234849, 2009.
- [7] A.M. Wazwaz. The modified decomposition method for analytic treatment of differential equations. *Appl. Math. Comput.* 2006, pp. 165-176.
- [8] G. Adomian. *Solving Frontier Problems of Physics: The Decomposition Method*. Boston: Kluwer Acad. Publ. 1994.
- [9] G.C. Wu, J.H. He. Fractional calculus of variations in fractal spacetime. *Nonlinear Science Letters*. 2010, **A 1**: 281–287.
- [10] J.H. He. Variational iteration method—a kind of nonlinear analytical technique: some examples. *International Journal of Nonlinear Mechanics*. 1999, **34**: 699–708.
- [11] J.H. He, X.H. Wu. Variational iteration method: new development and applications. *Computers & Mathematics with Applications*. 2007, **54**: 881–894.
- [12] J.H. He, G.C. Wu, F. Austin. The variational iteration method which should be followed. *Nonlinear Science Letters*. 2009, **A 1**: 1–30.
- [13] L.A. Soltani, A. Shirzadi. A new modification of the variational iteration method. *Computers & Mathematics with Applications*. 2010, **59**: 2528–2535.
- [14] N. Faraz, Y. Khan, A. Yildirim. Analytical approach to two-dimensional viscous flow with a shrinking sheet via variational iteration algorithm-II. *Journal of King Saud University*. doi:10.1016/j.jksus.2010.06.010.
- [15] G.C. Wu, E.W.M. Lee. Fractional variational iteration method and its application. *Physics Letters A*. doi:10.1016/j.physleta.2010.04.034.
- [16] E. Hesameddini, H. Latifizadeh. Reconstruction of variational iteration algorithms using the Laplace transform. *International Journal of Nonlinear Sciences and Numerical Simulation*. 2009, **10**: 1377–1382.
- [17] C. Chun. Fourier-series-based variational iteration method for a reliable treatment of heat equations with variable coefficients. *International Journal of Nonlinear Sciences and Numerical Simulation*. 2009, **10**: 1383–1388.
- [18] A.M. Wazwas. A study on linear and non-linear Schrödinger equations by the variational iteration method. *Chaos, Solitons and Fractals*. in press. doi:10.1016/j.chaos. 2006.10.009.
- [19] B. Jazbi and M. Moini. Application of He's homotopy perturbation method for Schrödinger equation. *Iranian*

- Journal of Mathematical Sciences and Informatics*. 2008, **3**(2): 13-19.
- [20] A. Ghorbani, J. Saberi-Nadjafi. He's homotopy perturbation method for calculating adomian polynomials. *International Journal of Nonlinear Sciences and Numerical Simulation*. 2007, **8**: 229–232.
- [21] A. Ghorbani. Beyond adomian's polynomials: He polynomials. *Chaos Solitons Fractals*. 2009, **39**: 1486–1492.
- [22] S.T. Mohyud-Din, M.A. Noor, K.I. Noor. Traveling wave solutions of seventh-order generalized KdV equation using He's polynomials. *International Journal of Nonlinear Sciences and Numerical Simulation*. 2009, **10**: 227–233.
- [23] Y. Khan, Q. Wu. Homotopy perturbation transform method for nonlinear equations using He's polynomials. *Computer and Mathematics with Applications*. doi:10.1016/j.camwa.2010.08.022.
- [24] S.T. Mohyud-Din, A. Yildirim. Homotopy perturbation method for advection problems. *Nonlinear Science Letters*. 2010, **A 1**: 307–312.
- [25] J.H. He. Homotopy perturbation method: a new nonlinear analytical technique. *Applied Mathematics and Computation*. 2003, **135**: 73–79.
- [26] J.H. He. Comparison of homotopy perturbation method and homotopy analysis method. *Applied Mathematics and Computation*. 2004, **156**: 527–539.
- [27] J.H. He. The homotopy perturbation method for nonlinear oscillators with discontinuities. *Applied Mathematics and Computation*. 2004, **151**: 287–292.
- [28] J.H. He. Homotopy perturbation method for bifurcation of nonlinear problems. *International Journal of Nonlinear Sciences and Numerical Simulation*. 2005, **6**: 207–208.
- [29] J.H. He. Some asymptotic methods for strongly nonlinear equation. *International Journal of Modern Physics*. 2006, **20**: 1144–1199.
- [30] J.H. He. Homotopy perturbation method for solving boundary value problems. *Physics Letters*. 2006, **A 350**: 87–88.
- [31] M. Rafei, D.D. Ganji. Explicit solutions of helmhotz equation and fifth-order KdV equation using homotopy perturbation method. *International Journal of Nonlinear Sciences and Numerical Simulation*. 2006, **7**: 321–328.
- [32] A.M. Siddiqui, R. Mahmood, Q.K. Ghori. Thin film flow of a third grade fluid on a moving belt by He's homotopy perturbation method. *International Journal of Nonlinear Sciences and Numerical Simulation*. 2006, **7**: 7–14.
- [33] D.D. Ganji. The applications of He's homotopy perturbation method to nonlinear equation arising in heat transfer. *Physics Letters*. 2006, **A 335**: 337–341.
- [34] L. Xu. He's homotopy perturbation method for a boundary layer equation in unbounded domain. *Computers & Mathematics with Applications*. 2007, **54**: 1067–1070.
- [35] J.H. He. An elementary introduction of recently developed asymptotic methods and nanomechanics in textile engineering. *International Journal of Modern Physics*. 2008, **22**: 3487–4578.
- [36] J.H. He. Recent developments of the homotopy perturbation method. *Topological Methods in Nonlinear Analysis*. 2008, **31**: 205–209.
- [37] E. Hesameddini, H. Latifzadeh. An optimal choice of initial solutions in the homotopy perturbation method. *International Journal of Nonlinear Sciences and Numerical Simulation*. 2009, **10**: 1389–1398.
- [38] E. Hesameddini, H. Latifzadeh. A new vision of the He's homotopy perturbation method. *International Journal of Nonlinear Sciences and Numerical Simulation*. 2009, **10**: 1415–1424.
- [39] J. Biazar, M. Gholami Porshokuhi, B. Ghanbari. Extracting a general iterative method from an adomian decomposition method and comparing it to the variational iteration method. *Computers & Mathematics with Applications*. 2010, **59**: 622–628.
- [40] S.A. Khuri. A Laplace decomposition algorithm applied to a class of nonlinear differential equations. *Journal of Applied Mathematics*. 2001, **1**: 141–155.
- [41] E. Yusufoglu. Numerical solution of Duffing equation by the Laplace decomposition algorithm. *Applied Mathematics and Computation*. 2006, **177**: 572–580.
- [42] Yasir Khan. An effective modification of the Laplace decomposition method for nonlinear equations. *International Journal of Nonlinear Sciences and Numerical Simulation*. 2009, **10**: 1373–1376.
- [43] Yasir Khan, Naeem Faraz. A new approach to differential difference equations. *Journal of Advanced Research in Differential Equations*. 2010, **2**: 1–12.
- [44] S. Islam, Y. Khan, N. Faraz, F. Austin. Numerical solution of logistic differential equations by using the Laplace decomposition method. *World Applied Sciences Journal*. 2010, **8**: 1100–1105.
- [45] M. Madani, M. Fathizadeh. Homotopy perturbation algorithm using Laplace transformation. *Nonlinear Science Letters*. 2010, **A 1**: 263–267.

- [46] S.T. Mohyud-Din, M.A. Noor and K.I. Noor. Traveling wave solutions of seventh-order generalized KdV equation using He's polynomials. *International Journal of Nonlinear Sciences and Numerical Simulation*. 2009, **10**: 227–233.
- [47] A. Yildirim. He's homotopy perturbation method for solving the space- and time- fractional telegraph equations. *International Journal of Computer Mathematics*. 2010, **87**(13): 2998-3006.
- [48] S.T. Mohyud-Din, A. Yildirim. Variational iteration method for solving Klein-Gordon equations. *Journal of Applied Mathematics, Statistics and Informatics*. 2010, **6**: 99-106.