

# Existence and Uniqueness of the Solution for Hilfer Neural Networks with Delays\*

Lulu Wang<sup>1</sup>, Yu Tian<sup>2,†</sup>

**Abstract** This paper mainly concerns with a class of nonlinear Hilfer fractional neutral recurrent neural networks with time varying delays. The existence and uniqueness of solutions in the space of weighted continuous functions are established by Banach's contraction principle. Finally, an example is provided to illustrate the application of the obtained results.

**Keywords** Hilfer fractional differential equation, existence and uniqueness, weighted space of continuous functions, fixed point theory

**MSC(2010)** 34A08, 34A12.

## 1. Introduction

In the past three decades, fractional calculus has received an increasing attention due to its various applications in engineering, mechanics, signal processing, material sciences, etc [1–4]. The related theories about fractional differential equations have been extensively studied by many researchers [5–8]. Additionally, Hilfer fractional derivative, which includes Riemann-Liouville fractional derivative and Caputo fractional derivative, was introduced in [1]. After that, a large number of fractional differential equations with Hilfer fractional derivatives were studied [9–13].

On the other hand, neutral neural networks with time delay have aroused naturally in a wide variety of fields like physics, chemistry, control, viscoelastic mechanics, porous media, electromagnetic and polymer rheology, etc. Therefore the issue concerning the existence, uniqueness of solutions of neutral neural networks has been widely discussed by many authors [14–19].

The successive approximation method [20, 21] and the fixed point theorems [20–30] have long been viewed as the main classical methods of studying existence and uniqueness problems in many areas of differential equations. Additionally, compared with fractional differential equations with delays, many dynamical systems not only depend on present and past states but also involve derivatives with delays, and neutral fractional differential equations with delays are often used to describe such systems [23, 31, 32]. Few authors studied the existence and uniqueness for a problem involving Hilfer fractional derivative. For example, C. Kou et

<sup>†</sup>the corresponding author.

Email address: [tianyu2992@bupt.edu.cn](mailto:tianyu2992@bupt.edu.cn) (Yu Tian), [wanglulu2432@163.com](mailto:wanglulu2432@163.com) (Lulu Wang)

<sup>1,2</sup>School of science, Beijing University of Posts and Telecommunications, Beijing 100876, China

\*The authors were supported by Education and teaching reform project of Beijing University of Posts and Telecommunications 2022SZ-A16 and Beijing University of Post and Telecommunications Graduate education and teaching reform and research project 2022Y026.

al. studied the existence and uniqueness on fractional differential equation with Riemann-Liouville fractional derivative [33]. D.F. Luo et.al proved the uniqueness of the solution for the stochastic fractional delay system with Caputo fractional derivative, see [31]. K.M. Furtati et al. studied the existence and uniqueness of global solutions for a class of nonlinear fractional differential equations involving Hilfer fractional derivative, see [34]. However, to the best of our knowledge, the existence and uniqueness of nonlinear Hilfer fractional neutral recurrent neural networks have not been yet developed. In this paper, we will study the existence and uniqueness of a class of nonlinear Hilfer fractional neutral recurrent neural networks with time varying delays by using Banach's fixed point theory.

In this paper, we consider a general class of neural networks with discrete and distributed varying delays which is described by

$$D_{0+}^{\alpha,\beta} \left[ x_i(t) - \sum_{j=1}^n q_{ij} x_j(t - \tau(t)) \right] = \sum_{j=1}^n a_{ij} f_j(x_j(t)) + \sum_{j=1}^n b_{ij} g_j(x_j(t - \tau(t))) + \sum_{j=1}^n l_{ij} \int_{t-r(t)}^t h_j(x_j(s)) ds, t \in [0, +\infty) \quad (1.1)$$

or

$$D_{0+}^{\alpha,\beta} \left[ x(t) - Qx(t - \tau(t)) \right] = Af(x(t)) + Bg(x(t - \tau(t))) + W \int_{t-r(t)}^t h(x(s)) ds,$$

where  $D_{0+}^{\alpha,\beta}$  is the Hilfer fractional derivative with  $0 < \alpha < 1$  and  $0 \leq \beta \leq 1$ , for  $i = 1, 2, 3, \dots, n$ ,  $x(t) = (x_1(t), x_2(t), \dots, x_n(t))^T \in \mathbb{R}^n$ ,  $(x_i(t) \in C([\vartheta, \infty), \mathbb{R}))$ , are the status vector relating to the neurons;  $A = (a_{ij})_{n \times n}$ ,  $B = (b_{ij})_{n \times n}$  and  $W = (l_{ij})_{n \times n}$  represent the connection weight matrix, delayed connection weight matrix and distributed delayed connection weight matrix, respectively;  $f_j, g_j, h_j$  are activation functions,  $f(x(t)) = (f_1(x_1(t)), f_2(x_2(t)), \dots, f_n(x_n(t)))^T \in \mathbb{R}^n$ ,  $g(x(t)) = (g_1(x_1(t)), g_2(x_2(t)), \dots, g_n(x_n(t)))^T \in \mathbb{R}^n$ ,  $h(x(t)) = (h_1(x_1(t)), h_2(x_2(t)), \dots, h_n(x_n(t)))^T \in \mathbb{R}^n$ , and the mappings  $f_j(\cdot), g_j(\cdot)$ , and  $h_j(\cdot)$  are globally Lipschitz continuous with constants  $\alpha_j, \beta_j$  and  $\gamma_j > 0$ , which satisfy  $f(0) \equiv 0, g(0) \equiv 0, h(0) \equiv 0$ , for  $j = 1, 2, \dots, n$ . Here  $\tau(t)$  and  $r(t)$  are nonnegative continuous functions that express discrete time varying delay and distributed time varying delay, respectively. Besides, the delays satisfy  $\lim_{t \rightarrow \infty} t - \tau(t) \rightarrow \infty$  and  $\lim_{t \rightarrow \infty} t - r(t) \rightarrow \infty$ . The initial condition for the system (1) is given by

$$I_{0+}^{(1-\alpha)(1-\beta)} \left[ x_i(0) - \sum_{j=1}^n q_{ij} x_j(-\tau(0)) \right] = \phi_i(0) - \sum_{j=1}^n q_{ij} \phi_j(-\tau(0)), \quad (1.2)$$

$$x(t) = \phi(t), t \in [\vartheta, 0], \quad (1.3)$$

where  $I_{0+}^{(1-\alpha)(1-\beta)}$  is the Riemann-Liouville fractional integral operator. Denote  $\vartheta = \inf_{t > 0} \{t - \tau(t), t - r(t)\}$ ,  $t \mapsto \phi(t) = (\phi_1(t), \phi_2(t), \dots, \phi_n(t))^T \in C([\vartheta, 0], L_{\mathcal{F}_0}^p(\Omega; \mathbb{R}^n))$ .

The purpose of this paper is to investigate the existence and uniqueness of the solution to the neutral delayed neural networks (1.1) with initial conditions (1.2)-(1.3) through fixed point method. The paper is organized as follows. Some necessary concepts and related lemmas are reviewed in Section 2. In Section 3, we prove the existence and uniqueness of the solution. Examples to illustrate our main results are given in Section 4.

## 2. Preliminaries and related lemmas

In this section, some definitions and lemmas are given which will be used throughout this paper.

**Definition 2.1.** ([1]) The right-sided Riemann-Liouville fractional integral of order  $\alpha > 0$ ,  $\alpha \in \mathbb{R}$  of a locally integrable function  $f$  is defined as

$$(I_{a^+}^\alpha f)(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-y)^{\alpha-1} f(y) dy,$$

for  $x > a$ , where  $\Gamma(\cdot)$  is the Gamma function.

**Definition 2.2.** ([1]) The right-sided fractional derivative of order  $0 < \alpha < 1$  and type  $0 \leq \beta \leq 1$  with respect to  $x$  is defined by

$$D_{a^+}^{\alpha, \beta} f(x) = \left( I_{a^+}^{\beta(1-\alpha)} \frac{d}{dx} (I_{a^+}^{(1-\beta)(1-\alpha)} f) \right)(x),$$

for functions  $f$  for which the expression on the right hand side exists, the special cases are Riemann-Liouville fractional derivative for  $\beta = 0$  and Caputo fractional difference for  $\beta = 1$ .

Different choices of norms can be considered. The norms we choose should be such that the space under consideration is complete and the equation yields a contraction mapping with respect to the norm. For the system (1.1) with initial conditions (1.2)-(1.3), we consider the weighted spaces of continuous functions

$$C_\gamma[0, b] = \{f : (0, b] \rightarrow \mathbb{R} : x^\gamma f(x) \in C[a, b]\}, 0 \leq \gamma < 1,$$

$$C_{1-\gamma}^{\alpha, \beta}[0, b] = \left\{ f \in C_{1-\gamma}[0, b], D_{0^+}^{\alpha, \beta} f \in C_{1-\gamma}[0, b] \right\},$$

and

$$C_{1-\gamma}^\gamma[0, b] = \left\{ f \in C_{1-\gamma}[0, b], D_{0^+}^\gamma f \in C_{1-\gamma}[0, b] \right\},$$

with the norms

$$\|f\|_{C_\gamma} = \|x^\gamma f(x)\|_C.$$

**Lemma 2.1** ([2]). Let  $0 \leq \gamma < 1$ ,  $0 < b < a$ ,  $g \in C_\gamma[0, b]$ ,  $g \in C[b, a]$  and  $g$  be continuous at  $b$ . Then  $g \in C_\gamma[0, a]$ .

The following lemmas provide some mapping properties of  $I_{0^+}^\alpha$ . Proofs can be found in [33].

**Lemma 2.2** ([2]). For  $x > 0$  we have

$$[I_{0^+}^\alpha t^{\beta-1}](x) = \frac{\Gamma(\beta)}{\Gamma(\beta + \alpha)} x^{\beta + \alpha - 1}, \alpha \geq 0, \beta > 0.$$

**Lemma 2.3** ([2]). For  $\alpha > 0$ , the fractional integration operator  $I_{0^+}^\alpha$  maps  $C[0, b]$  into  $C[0, b]$ .

**Lemma 2.4** ([2]). For  $\alpha > 0$  and  $0 \leq \gamma < 1$ , then the fractional integration operator  $I_{0^+}^\alpha$  is bounded from  $C_\gamma[0, b]$  into  $C_\gamma[0, b]$ .

**Lemma 2.5** ([34]). Let  $f \in L^1(a, c)$ . Then

$$\lim_{x \rightarrow c^+} \int_a^x (x-t)^{\alpha-1} f(t) dt = \int_a^c (c-t)^{\alpha-1} f(t) dt = \Gamma(\alpha) I_{a^+}^\alpha f(c), \alpha > 0.$$

### 3. Main result

In this section, we establish the existence of a unique solution to the problem (1.1)-(1.3) in the space  $C_{1-\gamma}^{\alpha, \beta}[0, b]$  by reducing the problem to a Volterra integral equation and then applying the Banach fixed point theorem. We start with some preparations.

**Lemma 3.1** (Theorem 23, [34]). Let  $\gamma = \alpha + \beta - \alpha\beta$  where  $0 < \alpha < 1$  and  $0 \leq \beta \leq 1$ . Let  $f : (a, b] \times \mathbb{R} \rightarrow \mathbb{R}$  be a function such that  $f(\cdot, y(\cdot)) \in C_{1-\gamma}[a, b]$  for any  $y \in C_{1-\gamma}[a, b]$ . If  $y \in C_{1-\gamma}^\alpha[a, b]$ , then  $y$  satisfies

$$\begin{cases} D_{a^+}^{\alpha, \beta} y(x) = f(x, y), x > a, 0 < \alpha < 1, 0 \leq \beta \leq 1, \\ I_{a^+}^{1-\gamma} y(a^+) = y_a, \end{cases}$$

if and only if  $y$  satisfies

$$y(x) = \frac{y_a}{\Gamma(\alpha + \beta - \alpha\beta)} (x-a)^{(\alpha-1)(1-\beta)} + \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t, y(t)) dt.$$

According to Lemma 3.1, one can easily express the equivalent Volterra integral equation of the system (1.1)-(1.3) in the following form

$$\begin{aligned} x(t) &= \frac{\left[ \phi_i(0) - \sum_{j=1}^n q_{ij} \phi_j(-\tau(0)) \right]}{\Gamma(\alpha + \beta - \alpha\beta)} t^{(\alpha-1)(1-\beta)} + \sum_{j=1}^n q_{ij} x_j(t - \tau(t)) \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \sum_{j=1}^n a_{ij} f_j(x_j(s)) ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \sum_{j=1}^n b_{ij} g_j(x_j(s - \tau(s))) ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \sum_{j=1}^n l_{ij} \int_{s-\tau(s)}^s h_j(x_j(u)) du ds. \end{aligned} \quad (3.1)$$

**Theorem 3.1.** Let  $0 \leq \alpha \leq 1$ ,  $0 < \beta < 1$ , and  $\gamma = \alpha + \beta - \alpha\beta$ . Let  $f, g, h : (0, a] \times \mathbb{R} \rightarrow \mathbb{R}$  be functions such that  $f(x(\cdot)), g(x(\cdot)), h(x(\cdot)) \in C_{1-\gamma}^{\beta(1-\alpha)}[0, a]$  for any  $x \in C_{1-\gamma}[0, a]$ . Besides,  $f(x(\cdot)), g(x(\cdot)), h(x(\cdot))$  are globally Lipschitz continuous with constants  $\alpha_j, \beta_j$  and  $\gamma_j > 0$ , satisfying that  $f(0) \equiv 0, g(0) \equiv 0, h(0) \equiv 0$ , for  $j = 1, 2, \dots, n$ , respectively. Then there exists a unique solution  $x(t)$  to the problem (1.1)-(1.3) such that  $x(t) - \sum_{j=1}^n q_{ij} x_j(t - \tau(t))$  in the space  $C_{1-\gamma}^\alpha[0, a]$ .

**Proof.** Define an operator  $Q$  by

$$\begin{aligned} Qx(t) &= \frac{[\phi_i(0) - \sum_{j=1}^n q_{ij}\phi_j(-\tau(0))]}{\Gamma(\alpha + \beta - \alpha\beta)} t^{(\alpha-1)(1-\beta)} + \sum_{j=1}^n q_{ij}x_j(t - \tau(t)) \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \sum_{j=1}^n a_{ij}f_j(x_j(s))ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \sum_{j=1}^n b_{ij}g_j(x_j(s - \tau(s)))ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \sum_{j=1}^n l_{ij} \int_{s-r(s)}^s h_j(x_j(u))duds, t \geq 0. \end{aligned} \quad (3.2)$$

First we prove the existence of a unique solution  $x(t)$  in the space  $C_{1-\gamma}[0, a]$ . Our proof is based on partitioning the interval  $(0, a]$  into subintervals on which the operator  $Q$  is a contraction, and based on the Banach's fixed point theorem. Note that  $C_{1-\gamma}[b_1, b_2]$ ,  $0 \leq b_1 < b_2 \leq a$  is a complete metric space with the metric  $d$  defined by

$$d(x_1, x_2) = \|x_1 - x_2\|_{C_{1-\gamma}[b_1, b_2]} := \max_{t \in [b_1, b_2]} |t^{1-\gamma}[x_1(t) - x_2(t)]|.$$

Choose  $t_1 \in (0, a]$  such that

$$\begin{aligned} \omega_1 &= \left( \sum_{j=1}^n |q_{ij}| \max_{t \in [0, t_1]} \left| 1 + \frac{\tau(t)}{t - \tau(t)} \right|^{1-\gamma} + \left[ \sum_{j=1}^n |a_{ij}| + \sum_{j=1}^n |b_{ij}| \max_{t \in [0, t_1]} \left| 1 + \frac{\tau(t)}{t - \tau(t)} \right|^{1-\gamma} \right. \right. \\ &\quad \left. \left. + \sum_{j=1}^n \frac{|l_{ij}|}{\gamma} \left( 1 - \max_{t \in [0, t_1]} \left| 1 - \frac{r(t)}{t} \right|^\gamma \right) \right] \frac{\Gamma(\gamma)}{\Gamma(\gamma + \alpha)} t_1^\alpha \right) < 1. \end{aligned} \quad (3.3)$$

Clearly, by Lemma 2.4, we obtain that  $Qx(t) \in C_{1-\gamma}[0, t_1]$ . Therefore  $Q : C_{1-\gamma}[0, t_1] \rightarrow C_{1-\gamma}[0, t_1]$ . Next we prove that  $Q$  is a contraction mapping. For any  $\varphi, \psi \in C_{1-\gamma}[0, t_1]$ , from Lemma 2.2, we have

$$\begin{aligned} &\| (Q\varphi)_i(t) - (Q\psi)_i(t) \|_{C_{1-\gamma}[0, t_1]} \\ &\leq \sum_{j=1}^n |q_{ij}| \max_{t \in [0, t_1]} \left| \frac{t^{1-\gamma}}{(t - \tau(t))^{1-\gamma}} (t - \tau(t))^{1-\gamma} (\varphi_j(t - \tau(t)) - \psi_j(t - \tau(t))) \right| \\ &\quad + \max_{t \in [0, t_1]} \left| t^{1-\gamma} \frac{1}{\Gamma(\alpha)} \sum_{j=1}^n a_{ij} \int_0^t \frac{(t-s)^{\alpha-1}}{s^{1-\gamma}} s^{1-\gamma} (f_j(\varphi_j(s)) - f_j(\psi_j(s))) ds \right| \\ &\quad + \max_{t \in [0, t_1]} \left| t^{1-\gamma} \frac{1}{\Gamma(\alpha)} \sum_{j=1}^n b_{ij} \int_0^t \frac{(t-s)^{\alpha-1}}{(s - \tau(s))^{1-\gamma}} (s - \tau(s))^{1-\gamma} \right. \\ &\quad \left. [g_j(\varphi_j(s - \tau(s))) - g_j(\psi_j(s - \tau(s)))] ds \right| \\ &\quad + \max_{t \in [0, t_1]} \left| t^{1-\gamma} \frac{1}{\Gamma(\alpha)} \sum_{j=1}^n l_{ij} \int_0^t (t-s)^{\alpha-1} \int_{s-r(s)}^s [h_j(\varphi_j(u)) - h_j(\psi_j(u))] duds \right| \end{aligned}$$

$$\begin{aligned}
&\leq \sum_{j=1}^n |q_{ij}| \max_{t \in [0, t_1]} \left| \frac{t^{1-\gamma}}{(t-\tau(t))^{1-\gamma}} \right| \|\varphi_j(t-\tau(t)) - \psi_j(t-\tau(t))\|_{C_{1-\gamma}[0, t_1]} \\
&\quad + \sum_{j=1}^n |a_{ij}| \frac{\Gamma(\gamma)}{\Gamma(\gamma+\alpha)} t_1^\alpha \|\varphi_j(t) - \psi_j(t)\|_{C_{1-\gamma}[0, t_1]} \\
&\quad + \sum_{j=1}^n |b_{ij}| \max_{t \in [0, t_1]} \left| 1 + \frac{\tau(t)}{t-\tau(t)} \right|^{1-\gamma} \frac{\Gamma(\gamma)}{\Gamma(\gamma+\alpha)} t_1^\alpha \|\varphi_j(t) - \psi_j(t)\|_{C_{1-\gamma}[0, t_1]} \\
&\quad + \sum_{j=1}^n \frac{|l_{ij}|}{\gamma} \left( 1 - \max_{t \in [0, t_1]} \left| 1 - \frac{r(t)}{t} \right|^\gamma \right) \frac{\Gamma(\gamma)}{\Gamma(\gamma+\alpha)} t_1^\alpha \|\varphi_j(t) - \psi_j(t)\|_{C_{1-\gamma}[0, t_1]} \\
&\leq \left( \sum_{j=1}^n |q_{ij}| \max_{t \in [0, t_1]} \left| 1 + \frac{\tau(t)}{t-\tau(t)} \right|^{1-\gamma} + \left[ \sum_{j=1}^n |a_{ij}| + \sum_{j=1}^n |b_{ij}| \max_{t \in [0, t_1]} \left| 1 + \frac{\tau(t)}{t-\tau(t)} \right|^{1-\gamma} \right. \right. \\
&\quad \left. \left. + \sum_{j=1}^n \frac{|l_{ij}|}{\gamma} \left( 1 - \max_{t \in [0, t_1]} \left| 1 - \frac{r(t)}{t} \right|^\gamma \right) \right] \frac{\Gamma(\gamma)}{\Gamma(\gamma+\alpha)} t_1^\alpha \right) \|\varphi_j(t) - \psi_j(t)\|_{C_{1-\gamma}[0, t_1]}.
\end{aligned}$$

Therefore,

$$\|(Q\varphi)_i(t) - (Q\psi)_i(t)\|_{C_{1-\gamma}[0, t_1]} \leq \omega_1 \|\varphi_j(t) - \psi_j(t)\|_{C_{1-\gamma}[0, t_1]}.$$

By the contraction mapping principle, we obtain that  $Q$  has a unique fixed point  $x_0^*(t) \in C_{1-\gamma}[0, t_1]$ .

If  $t_1 \neq a$ , then we consider the interval  $[t_1, a]$ . On this interval we consider solution  $x(t) \in C[t_1, a]$  to the equation

$$\begin{aligned}
x(t) = Qx(t) &:= \frac{\left[ \phi_i(0) - \sum_{j=1}^n q_{ij} \phi_j(-\tau(0)) \right]}{\Gamma(\alpha + \beta - \alpha\beta)} t^{(\alpha-1)(1-\beta)} + \sum_{j=1}^n q_{ij} x_j(t-\tau(t)) \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} (t-s)^{\alpha-1} \sum_{j=1}^n a_{ij} f_j(x_0^*(s)) ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_{t_1}^t (t-s)^{\alpha-1} \sum_{j=1}^n a_{ij} f_j(x_j(s)) ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} (t-s)^{\alpha-1} \sum_{j=1}^n b_{ij} g_j(x_0^*(s-\tau(s))) ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_{t_1}^t (t-s)^{\alpha-1} \sum_{j=1}^n b_{ij} g_j(x_j(s-\tau(s))) ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} (t-s)^{\alpha-1} \sum_{j=1}^n l_{ij} \int_{s-r(s)}^s h_j(x_0^*(u)) du ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_{t_1}^t (t-s)^{\alpha-1} \sum_{j=1}^n l_{ij} \int_{s-r(s)}^s h_j(x_j(u)) du ds. \tag{3.4}
\end{aligned}$$

Now we select  $t_2 \in [t_1, a]$  such that

$$\omega_2 = \sum_{j=1}^n |q_{ij}| + \frac{(t_2 - t_1)^\alpha}{\alpha \Gamma(\alpha)} \left( \sum_{j=1}^n |a_{ij} \alpha_j| + \sum_{j=1}^n |b_{ij} \beta_j| + r \sum_{j=1}^n |l_{ij} \gamma_j| \right) < 1. \quad (3.5)$$

Since  $x_i(t) \in C[t_1, t_2]$ , and  $f(x(t)), g(x(t)), h(x(t)) \in C[t_1, t_2]$  for any  $x(t) \in C[t_1, t_2]$ , Lemma 2.3 implies that the right-hand side of (3.4) is in  $C[t_1, t_2]$ . Therefore  $Q : C[t_1, t_2] \rightarrow C[t_1, t_2]$ . For any  $\varphi, \psi \in C[t_1, t_2]$ , we have

$$\begin{aligned} & \| (Q\varphi)_i(t) - (Q\psi)_i(t) \|_{C[t_1, t_2]} \\ & \leq \left\| \sum_{j=1}^n q_{ij} (\varphi_j(t - \tau(t)) - \psi_j(t - \tau(t))) \right\|_{C[t_1, t_2]} \\ & \quad + \left\| \frac{1}{\Gamma(\alpha)} \sum_{j=1}^n a_{ij} \int_{t_1}^t (t-s)^{\alpha-1} (f_j(\varphi_j(s)) - f_j(\psi_j(s))) ds \right\|_{C[t_1, t_2]} \\ & \quad + \left\| \frac{1}{\Gamma(\alpha)} \sum_{j=1}^n b_{ij} \int_{t_1}^t (t-s)^{\alpha-1} [g_j(\varphi_j(s - \tau(s))) - g_j(\psi_j(s - \tau(s)))] ds \right\|_{C[t_1, t_2]} \\ & \quad + \left\| \frac{1}{\Gamma(\alpha)} \sum_{j=1}^n l_{ij} \int_{t_1}^t (t-s)^{\alpha-1} \int_{s-r(s)}^s [h_j(\varphi_j(u)) - h_j(\psi_j(u))] du ds \right\|_{C[t_1, t_2]} \\ & \leq \sum_{j=1}^n |q_{ij}| \max \left\{ \max_{t \in [t_1 - \tau, t_1]} |\varphi_j(t) - \psi_j(t)|, \max_{t \in [t_1, t_2]} |\varphi_j(t) - \psi_j(t)| \right\} \\ & \quad + \frac{1}{\Gamma(\alpha)} \sum_{j=1}^n |a_{ij} \alpha_j| \left\| \max_{s \in [t_1, t]} (\varphi_j(s) - \psi_j(s)) \int_{t_1}^t (t-s)^{\alpha-1} ds \right\|_{C[t_1, t_2]} \\ & \quad + \frac{1}{\Gamma(\alpha)} \sum_{j=1}^n |b_{ij} \beta_j| \left\| \max_{s \in [t_1, t]} (\varphi_j(s - \tau(s)) - \psi_j(s - \tau(s))) \int_{t_1}^t (t-s)^{\alpha-1} ds \right\|_{C[t_1, t_2]} \\ & \quad + \frac{r}{\Gamma(\alpha)} \sum_{j=1}^n |l_{ij} \gamma_j| \left\| \max_{u \in [s-r(s), s]} (\varphi_j(u) - \psi_j(u)) \int_{t_1}^t (t-s)^{\alpha-1} ds \right\|_{C[t_1, t_2]} \\ & \leq \sum_{j=1}^n |q_{ij}| \|\varphi_j(t) - \psi_j(t)\|_{C[t_1, t_2]} + \frac{(t_2 - t_1)^\alpha}{\alpha \Gamma(\alpha)} \sum_{j=1}^n |a_{ij} \alpha_j| \|\varphi_j(t) - \psi_j(t)\|_{C[t_1, t_2]} \\ & \quad + \frac{(t_2 - t_1)^\alpha}{\alpha \Gamma(\alpha)} \sum_{j=1}^n |b_{ij} \beta_j| \|\varphi_j(t) - \psi_j(t)\|_{C[t_1, t_2]} \\ & \quad + \frac{r(t_2 - t_1)^\alpha}{\alpha \Gamma(\alpha)} \sum_{j=1}^n |l_{ij} \gamma_j| \|\varphi_j(t) - \psi_j(t)\|_{C[t_1, t_2]} \\ & = \left( \sum_{j=1}^n |q_{ij}| + \frac{(t_2 - t_1)^\alpha}{\alpha \Gamma(\alpha)} \left( \sum_{j=1}^n |a_{ij} \alpha_j| + \sum_{j=1}^n |b_{ij} \beta_j| + r \sum_{j=1}^n |l_{ij} \gamma_j| \right) \right) \\ & \quad \|\varphi_j(t) - \psi_j(t)\|_{C[t_1, t_2]}. \end{aligned}$$

Since  $0 < \omega_2 < 1$ ,  $Q$  is a contraction. By the contraction mapping principle, we obtain that  $Q$  has a unique fixed point  $x_1^*(t) \in C[t_1, t_2]$  to (3.2). Moreover, it follows

from Lemma 2.5 that  $x_0^*(t_1) = x_1^*(t_1)$ . Therefore, if

$$x^*(t) = \begin{cases} x_0^*(t), & 0 < t \leq t_1, \\ x_1^*(t), & t_1 < t \leq t_2, \end{cases}$$

then by Lemma 2.1,  $x^*(t) \in C_{1-\gamma}[0, t_2]$ . So  $x^*(t)$  is the unique solution of (3.2) in  $C_{1-\gamma}[0, t_2]$  on the interval  $[0, t_2]$ .

If  $t_2 \neq a$ , we repeat the process as necessary, such as  $N - 2$  times, to obtain the unique solution  $x_k^*(t) \in C[t_k, t_{k+1}]$ ,  $k = 2, 3, \dots, N$ , where  $0 = t_0 < t_1 < \dots < t_N = a$ , such that

$$\omega_{k+1} = \sum_{j=1}^n |q_{ij}| + \frac{(t_{k+1} - t_k)^\alpha}{\alpha \Gamma(\alpha)} \left( \sum_{j=1}^n |a_{ij} \alpha_j| + \sum_{j=1}^n |b_{ij} \beta_j| + r \sum_{j=1}^n |l_{ij} \gamma_j| \right) < 1.$$

As a result we have the unique solution  $x^*(t) \in C_{1-\gamma}[0, a]$  of (3.2) given by

$$x^*(t) = x_k^*(t), t \in (t_k, t_{k+1}], k = 0, 1, \dots, N - 1.$$

It remains to show that such a unique solution  $x^*(t) \in C_{1-\gamma}[0, a]$  is actually in  $C_{1-\gamma}^\gamma[0, a]$ . Applying  $D_{0+}^\gamma$  to both sides yields of (3.1)

$$\begin{aligned} & D_{0+}^\gamma \left[ x^*(t) - \sum_{j=1}^n q_{ij} x_j^*(t - \tau(t)) \right] \\ &= D_{0+}^\gamma \left[ I_{0+}^\alpha \left( \sum_{j=1}^n a_{ij} f_j(x_j(t)) + \sum_{j=1}^n b_{ij} g_j(x_j(t - \tau(t))) + \sum_{j=1}^n l_{ij} \int_{t-\tau(t)}^t h_j(x_j(s)) ds \right) \right] \\ &= D_{0+}^{\beta(1-\alpha)} \left[ \sum_{j=1}^n a_{ij} f_j(x_j(t)) + \sum_{j=1}^n b_{ij} g_j(x_j(t - \tau(t))) + \sum_{j=1}^n l_{ij} \int_{t-\tau(t)}^t h_j(x_j(s)) ds \right]. \end{aligned}$$

By hypothesis, the right hand side is in  $C_{1-\gamma}[0, a]$  and thus  $x^*(t) - \sum_{j=1}^n q_{ij} x_j^*(t - \tau(t)) \in C_{1-\gamma}^\gamma[0, a]$ .

Therefore, by Lemma 3.1,  $x^*(t)$  is the unique solution of (1.1)-(1.3) such that  $x^*(t) - \sum_{j=1}^n q_{ij} x_j^*(t - \tau(t))$  in the space  $C_{1-\gamma}^\gamma[0, a]$ .  $\square$

## 4. Example

**Example 4.1.** Consider the following two-dimensional Hilfer fractional neutral neural network

$$D_{0+}^{\alpha, \beta} \left[ x(t) - Qx(t - \tau(t)) \right] = Af(x(t)) + Bg(x(t - \tau(t))) + W \int_{t-\tau(t)}^t h(x(s)) ds, \quad (4.1)$$

where  $\alpha = 0.8, \beta = 0.25$ ,

$$Q = \begin{pmatrix} -0.1 & 0 \\ 0.1 & 0.1 \end{pmatrix}, A = \begin{pmatrix} 0.2 & 0.4 \\ 0.3 & 0.2 \end{pmatrix}, B = \begin{pmatrix} -0.1 & 0.2 \\ 0.1 & 0.2 \end{pmatrix}, W = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix},$$



$f(x) = g(x) = h(x) = 0.2 \tanh(x)$ .  $\tau(t), r(t)$  are continuous functions such that  $|\tau(t)| \leq 1$  and  $|r(t)| \leq 1$ .

In this example, Letting  $\alpha_j = 0.2, \beta_j = 0.25, \gamma_j = 0.1, j = 1, 2, k = 2, a = 1, t_1 = 0.1, t_2 = 0.2, t_3 = a = 1$ , we get

$$\omega_1 = \left( \sum_{j=1}^2 |q_{ij}| \max_{t \in [0, t_1]} \left| 1 + \frac{\tau(t)}{t - \tau(t)} \right|^{1-\gamma} + \left[ \sum_{j=1}^2 |a_{ij}| + \sum_{j=1}^2 |b_{ij}| \max_{t \in [0, t_1]} \left| 1 + \frac{\tau(t)}{t - \tau(t)} \right|^{1-\gamma} + \sum_{j=1}^2 \frac{|l_{ij}|}{\gamma} \left( 1 - \max_{t \in [0, t_1]} \left| 1 - \frac{r(t)}{t} \right|^\gamma \right) \right] \frac{\Gamma(\gamma)}{\Gamma(\gamma + \alpha)} t_1^\alpha \right) \leq 0.671 < 1,$$

$$\omega_2 = \sum_{j=1}^n |q_{ij}| + \frac{(t_2 - t_1)^\alpha}{\alpha \Gamma(\alpha)} \left( \sum_{j=1}^n |a_{ij} \alpha_j| + \sum_{j=1}^n |b_{ij} \beta_j| + r \sum_{j=1}^n |l_{ij} \gamma_j| \right) \leq 0.604 < 1.$$

Letting  $k = 2$ , we have

$$\omega_3 = \sum_{j=1}^2 |q_{ij}| + \frac{(t_{k+1} - t_k)^\alpha}{\alpha \Gamma(\alpha)} \left( \sum_{j=1}^2 |a_{ij} \alpha_j| + \sum_{j=1}^2 |b_{ij} \beta_j| + r \sum_{j=1}^2 |l_{ij} \gamma_j| \right) \leq 0.897 < 1.$$

Applying Theorem 3.1, the initial value problem (4.1) has a unique solution  $x(t)$  such that  $x(t) - \sum_{j=1}^n q_{ij} x_j(t - \tau(t))$  in the space  $C_{1-\gamma}^\gamma[0, a]$ .

## References

- [1] R. Hilfer, Applications of Fractional Calculus in Physics, World Scientific, Singapore, 2000.
- [2] A. Kilbas, H. Srivastava, J. Trujillo, Theory and applications of fractional differential equations, Elsevier Science Limited, Amsterdam, 2006.
- [3] J. Wang, M. Feckan and Y. Zhou, A survey on impulsive fractional differential equations, Fract. Calc. Appl. Anal. 19 (2016) 806-831.
- [4] Y. Zhou and L. Peng, On the time-fractional Navier-Stokes equations, Comput. Math. Appl. 73 (2017) 874-891.
- [5] J. Wang and Y. Zhang, Nonlocal initial value problems for differential equations with Hilfer fractional derivative, Appl. Math. Comput. 266 (2015) 850-859.
- [6] H. Gu and J. Trujillo, Existence of mild solution for evolution equation with Hilfer fractional derivative, Appl. Math. Comput. 257 (2015) 344-354.
- [7] A. Debbouche and V. Antonov, Approximate controllability of semilinear Hilfer fractional differential inclusions with impulsive control inclusion conditions in banach spaces, Chaos Solitons Fractals. 102 (2017) 140-148.
- [8] H. Ahmed and M. El-Borai, Hilfer fractional stochastic integro-differential equations, Appl. Math. Comput. 331 (2018) 182-189.
- [9] Syed Sabyel. Haider, Mujeeb ur. Rehman and Thabet. Abdeljawad, On Hilfer fractional difference operator, Adv. Difference Equations. 1 (2020) 1-20.
- [10] J. Lv and X. Yang, A class of Hilfer fractional stochastic differential equations and optimal controls, Adv. Difference Equations. 1 (2019) 1.

- [11] Joel. Restrepo and Durvudkhan. Suragan, Hilfer-type fractional differential equations with variable coefficients, *Chaos Solitons Fractals*. 150 (2021) 111146.
- [12] R. Hilfer, Experimental evidence for fractional time evolution in glass forming materials. *Chem Phys*. 284 (2002) 399-408.
- [13] D. Luo, Q. Zhu and Z. Luo, A novel result on averaging principle of stochastic Hilfer-type fractional system involving non-Lipschitz coefficients, *Appl. Math. Lett.* 122 (2021) 107549.
- [14] A. Djoudi and R. Khemis, Fixed points techniques and stability for neutral nonlinear differential equations with unbounded delays, *Georgian Math. J.* 13 (2006) 25-34.
- [15] S. Haykin, *Neural Networks*, Prentice-Hall, Englewood Cliffs, 1994.
- [16] M. Forti and A. Tesi, New conditions for global stability of neural networks with application to linear and quadratic programming problems, *IEEE Trans. Circuits Syst. I.* 42 (1995) 354-366.
- [17] X. Liao and J. Wang, Algebraic criteria for global exponential stability of cellular neural networks with multiple time delays, *IEEE Trans. Circuits Syst. I.* 50 (2003) 268-275.
- [18] Z. Zeng and D. Wang, Global stability of a general class of discrete-time recurrent neural networks, *Neural Comput.* 18 (2006) 848-870.
- [19] Z. Zeng and J. Wang, Complete stability of cellular neural networks with time-varying delays, *IEEE Trans. Circuits Syst. I.* 53 (2006) 944-955.
- [20] G. Chen, O. van Gaans and S. Lunel, Existence and exponential stability of a class of impulsive neutral stochastic partial differential equations with delays and poisson jumps, *Statist. Probab. Lett.* 141 (2018) 7-18.
- [21] F. Jiang and Y. Shen, A note on the existence and uniqueness of mild solutions to neutral stochastic partial functional differential equations with non-Lipschitz coefficients, *Comput. Math. Appl.* 61 (2011) 1590-1594.
- [22] J. You and S. Sun, Mixed Boundary Value Problems for a Class of Fractional Differential Equations with Impulses, *Journal of Nonlinear Modeling and Analysis*, 3 (2021) 263-273.
- [23] J. Luo and Takeshi. Taniguchi, Fixed points and stability of stochastic neutral partial differential equations with infinite delays, *Stochastic Anal. Appl.* 27 (2009) 1163-1173.
- [24] J. Zhu, X. Wang and X. Fu, Existence and Blowup of Solutions for Neutral Partial Integro-differential Equations with State-dependent Delay, *Journal of Nonlinear Modeling and Analysis*, 2 (2020) 287-313.
- [25] G. Chen, D. Li, L. Shi, O. van Gaans. Onno and S. Lunel, Stability results for stochastic delayed recurrent neural networks with discrete and distributed delays. *J. Differential Equations*. 264 (2018) 3864-3898.
- [26] B. Ahmad, S. Ntouyas and A. Alsaedi, An existence result for fractional differential inclusions with nonlinear integral boundary conditions, *J. Inequal. Appl.* 1 (2013) 296-404.
- [27] H. Baghani and J. Nieto, On fractional Langevin equation involving two fractional orders in different intervals, *Nonlinear Anal. Model. Control*, 24 (2019) 884-897.

- [28] H. Fazli and J. Nieto, Fractional Langevin equation with anti-periodic boundary conditions, *Chaos Soliton Fract.* 114 (2018) 332-337.
- [29] R. Ghorbanian, V. Hedayati, M. Postolachei, S. Rezaour, On a fractional differential inclusion via a new integral boundary condition, *J. Inequal. Appl.* 2014 (2014) 1-20.
- [30] H. Lian, P. Wang, W. Ge, Unbounded upper and lower solutions method for Sturm-Liouville boundary value problem on infinite intervals, *Nonlinear Anal.* 70 (2009) 2627-2633.
- [31] D. Luo, M. Tian and Q. Zhu, Some results on finite-time stability of stochastic fractional-order delay differential equations. *Chaos Solitons Fractals*, 158 (2022) 111996.
- [32] F. Jiang and Y. Shen, Stability of impulsive stochastic neutral partial differential equations with infinite delays, *ASIAN J CONTROL.* 14 (2012) 1706-1709.
- [33] C. Kou, J. Liu and Y. Ye, Existence and uniqueness of solutions for the Cauchy-type problems of fractional differential equations, *Discrete Dyn. Nat. Soc.* 2010 (2010) 1-15.
- [34] K. Furati, M. Kassim and N. Tatar, Existence and uniqueness for a problem involving Hilfer fractional derivative, *Comput. Math. Appl.* 64 (2012) 1616-1626.