

# Here is the Title

John Doe<sup>1,2</sup> and Jane Smith<sup>1,2,\*</sup>

<sup>1</sup>Department of Chemistry, Faculty of Arts and Sciences, Beijing Normal University, Zhuhai 519087, China

<sup>2</sup>College of Chemistry, Beijing Normal University, Beijing 100875, China

\*Corresponding authors: xxxxx@xxx

Received: 20 Oct. 2026; Revision: 25 Oct. 2026; Accepted: 25 Dec. 2026; Online: 28 Dec. 2026.

**Abstract:** The abstract should be a single paragraph that summarises the content of the article.


**Key words:** keyword1, keyword2, keyword3, keyword4.

The main text of the article should appear here. The references should be cited as[1–8].

The time-dependent Schrödinger equation is:

$$i\hbar \frac{\partial \Psi(\mathbf{r}, t)}{\partial t} = \hat{H}\Psi(\mathbf{r}, t). \quad (1)$$

All figures and tables must be mentioned in the text consecutively and numbered with Arabic numerals, such as Figure 1 or Table 1.



**Figure 1.** Each figure must have a caption that includes the figure number and a brief description, preferably one or two sentences.

**Table 1.** Each table must have a brief (one phrase or sentence) title that describes its contents.

Head 1 <sup>[a]</sup>	Head 2	Head 3	Head 4
Column 1	Column 2 <sup>[b]</sup>	Column 3	Column 4
Column 1	Column 2	Column 3	Column 4
Column 1	Column 2	Column 3	Column 4
Column 1	Column 2	Column 3	Column 4
Column 1	Column 2	Column 3	Column 4

[a] Table footnote. [b] Table footnote.

## Acknowledgments

Here is the place to acknowledge people, organizations, and financing (you may state grant numbers and sponsors here).

## Supporting information

The online version contains supplementary material available at website xxx. A listing of the contents of each file supplied as Supporting Information should be included.

## Notes

The authors declare no competing financial interest.

## References

- [1] Berger M. J. and Colella P., Local adaptive mesh refinement for shock hydrodynamics, *J. Comput. Phys.*, **82** (1989), 62–84.
- [2] Tan Z. J., Tang T. and Zhang Z. R., A simple moving mesh method for one- and two-dimensional phase-field equations, *J. Comput. Appl. Math.*, To appear.
- [3] McKechnie S., Frost J. M., Pashov D., Azarhoosh P., Walsh A. and Schilfgaarde M., Dynamic symmetry breaking and spin splitting in metal halide perovskites, 2017, DOI: 10.48550/arXiv.1711.00533.
- [4] de Boor C., Good approximation by splines with variable knots II, In *Springer Lecture Notes Series 363*, Springer-Verlag, Berlin, 1973.
- [5] Tullius T. D., Supramolecular reactivity and transport: Bioinorganic systems, In *Comprehensive Supramolecular Chemistry*, Vol. **5**, (Eds. Atwood J. L., Davies J. E. D., MacNicol D. D., Vögtle F. and Suslick K. S.), Pergamon, Oxford, 1996, pp. 317–343.
- [6] Doe J., *Dataset title*, Year, DOI: persistent identifier (e.g. DOI).
- [7] Sheldrick G. M., *SHELXS-96, program for the solution of crystal structures*, University of Göttingen, Göttingen, Germany, 1996.
- [8] Tullius T. D., *Title of PhD thesis*, University of Newcastle, UK, 1991.