

Blowing-Up Solutions of the Shallow Water Equations*

Nurbol Koshkarbayev[†]

Abstract In the paper, we study the question about global unsolvability of the Kawahara and Kaup-Kupershmidt shallow water equations in a bounded domain. For certain initial-boundary-value problems of the shallow-water equations, we establish the necessary conditions for the existence of global solutions. The proof of the results is based on the nonlinear capacity method. In closing, we provide some examples.

Keywords Kaup-Kupershmidt equation, Kawahara equation, shallow water, blow-up solution, mixed problem

MSC(2010) 35K55, 35R11.

1. Introduction

In the paper, we consider some shallow water equations as follows

$$\partial_t u + \alpha \partial_x^5 u + \beta \partial_x^3 u + \gamma \partial_x u + u \partial_x u = 0, (x, t) \in (a, b) \times (0, T), \quad (1.1)$$

$$\partial_t u + \alpha \partial_x^5 u + \beta \partial_x^3 u + \gamma \partial_x u - \partial_x u \partial_x^2 u = 0, (x, t) \in (a, b) \times (0, T), \quad (1.2)$$

with the initial data

$$u(x, 0) = u_0(x), \quad (1.3)$$

where $-\infty < a < b < +\infty$, $\alpha \neq 0$, β, γ are real numbers.

The model (1.1) is also called the Kawahara equation [9]. It arises in the study of the water waves with surface tension, in which the Bond number takes on the critical value, where the Bond number represents a dimensionless magnitude of surface tension in the shallow water regime (see [3, 10]). Equation (1.2) is often called the Kaup-Kupershmidt equation [8]. This model has arisen from the study of the capillary-gravity waves [2, 6, 7, 21].

If $\alpha = 0$, then the model (1.1) reduces to the well-known Korteweg-de Vries equation [4]

$$\partial_t u + \beta \partial_x^3 u + \gamma \partial_x u + u \partial_x u.$$

The local and global well-posedness of problems (1.1) - (1.3) for $\alpha = -1$, $\beta = 1$, $\gamma = 0$, with the boundary conditions

$$u(0, t) = 0, u(1, t) = 0, t \geq 0,$$

[†]the corresponding author.

Email address: koshkarbaev.nurbol@gmail.com (N. Koshkarbayev)
Institute of Mathematics and Mathematical Modeling, 125 Pushkin str.,
050010 Almaty, Kazakhstan

*This research has been funded by the Science Committee of the Ministry of Higher Education and Science of the Republic of Kazakhstan Grant No. AP22686595.

$$\begin{aligned}\partial_x u(0, t) &= 0, \quad \partial_x u(1, t) = 0, \quad t \geq 0, \\ \partial_x^2 u(1, t) &= 0, \quad t \geq 0,\end{aligned}$$

were studied by Larkin et al. in [12–14]. Note that the local well-posedness of the equations (1.1)–(1.2) with the initial data (1.3) and several boundary conditions was investigated in [1, 5, 15, 18, 20, 22, 23].

The main aim of this paper is to obtain the blowing-up solutions of the above equations, more precisely, solutions that blow up in finite time for the large class of boundary conditions. Our approach is based on the Mitidieri-Pohozaev nonlinear capacity method (see [16, 17], also [11, 19]), more precisely, on the choice of specific functions corresponding to the initial and boundary conditions.

A small difference in our approach is that we will study shallow-water equations without boundary conditions. We suppose the boundary conditions are such that there exists a sufficiently smooth function φ , for which the functional $B(u, \phi)$, containing u and φ and their k -derivatives, is lower bounds by a certain functional of φ . This allows us to study some classes of boundary conditions.

2. Blow-up of solution of the Kawahara equation

Let us consider the function $\varphi \in C^5([a, b])$ defined on the domain $a < x < b$ with arbitrary parameters $a, b \in \mathbb{R}$ and monotonically nondecreasing:

$$\varphi'(x) \geq 0 \quad \text{for } x \in [a, b], \quad (2.1)$$

and let φ satisfy the following properties:

$$\begin{cases} \theta_1 := \int_a^b \frac{(\alpha\varphi^{(5)} + \beta\varphi''' + \gamma\varphi')^2}{\varphi'} dx < \infty; \\ \theta_2 := \int_a^b \frac{\varphi^2}{\varphi'} dx < \infty. \end{cases} \quad (2.2)$$

Suppose that there is a classical solution $u(x, t) \in C_{t,x}^{1,5}(\mathbb{R} \times (0, T))$.

Multiplying Kawahara shallow water equation (1.1) by φ , we have

$$\begin{aligned}\int_a^b \partial_t u(x, t) \varphi(x) dx &= -\alpha \int_a^b \partial_x^5 u(x, t) \varphi(x) dx - \beta \int_a^b \partial_x^3 u(x, t) \varphi(x) dx \\ &\quad - \gamma \int_a^b \partial_x u(x, t) \varphi(x) dx - \int_a^b u(x, t) \partial_x u(x, t) \varphi(x) dx.\end{aligned}$$

Applying integration by parts, we arrive at

$$\begin{aligned}\partial_t \int_a^b u(x, t) \varphi(x) dx &= \alpha \int_a^b u(x, t) \varphi^{(5)}(x) dx + \beta \int_a^b u(x, t) \varphi'''(x) dx \\ &\quad + \gamma \int_a^b u(x, t) \varphi'(x) dx + \frac{1}{2} \int_a^b u^2(x, t) \varphi'(x) dx \\ &\quad + \mathcal{B}(u(x, t), \varphi(x)) \Big|_{x=a}^{x=b},\end{aligned} \quad (2.3)$$