

## CONVERGENCE AND SUPERCONVERGENCE OF HERMITE BICUBIC ELEMENT FOR EIGENVALUE PROBLEM OF THE BIHARMONIC EQUATION\*

Dong-sheng Wu

*(Institute of Systems Science, Academy of Mathematics and Systems Sciences, Chinese Academy of Sciences, Beijing 100080, China )*

### Abstract

In this paper, we discuss the convergence and superconvergence for eigenvalue problem of the biharmonic equation by using the Hermite bicubic element. Based on asymptotic error expansions and interpolation postprocessing, we gain the following estimation:

$$0 \leq \bar{\lambda}_h - \lambda \leq C_\epsilon h^{8-\epsilon}$$

where  $\epsilon > 0$  is an arbitrary small positive number and  $C_\epsilon > 0$  is a constant.

*Key words:* Hermite bicubic element, Biharmonic equation, Interpolation postprocessing, Eigenvalue problem.

Eigenvalue problem for biharmonic equation is an interesting and important problem, see Ciarlet and Lions<sup>[3]</sup>. In 1979, Rannacher<sup>[8]</sup> used the Adini nonconforming finite element to solve this problem and obtained:

$$|\lambda_h - \lambda| \leq Ch^2.$$

Recently, Yang<sup>[6]</sup> has proved that the order of convergence of  $\lambda_h$  is just 2. The aim of this paper is to improve the order of convergence by using Hermite bicubic element. To our knowledge, there is not any result for approximation to the eigenvalue problem by using this element in literature. This paper fills in this gap. Not only the convergence is discussed:

$$0 \leq \lambda_h - \lambda \leq Ch^4.$$

but also a superconvergence is presented:

$$0 \leq \bar{\lambda}_h - \lambda \leq C_\epsilon h^{8-\epsilon}.$$

where  $\epsilon > 0$  is an arbitrary small positive number.

Note that the degree of freedom is 12 for Adini element and 16 for Hermite bicubic element, respectively. We can see that the increase of the degree of freedom for Hermite element is not too much, but the order of convergence and superconvergence by using Hermite element is one times and even two times higher than that of Adini element scheme, respectively. We mention that our techniques here are based on asymptotic error expansions and interpolation postprocessing<sup>[1,2,5,7]</sup>.

Let  $\Omega$  be a rectangle domain. We will consider the following eigenvalue problem of biharmonic equation :

$$\begin{cases} \Delta^2 u = \lambda u & \text{in } \Omega \\ u|_{\partial\Omega} = \frac{\partial u}{\partial n}|_{\partial\Omega} = 0 \end{cases}$$

Its weak form reads as: find a pair of  $(\lambda, u)$ ,  $\lambda \in R$ ,  $u \in H_0^2(\Omega)$ ,  $\|u\|_0 = 1$  such that

$$a(u, v) = \lambda(u, v), \quad \forall v \in H_0^2(\Omega), \tag{1}$$

---

\* Received December 2, 1997.

where

$$a(u, v) = \int_{\Omega} (u_{xx}v_{xx} + 2u_{xy}v_{xy} + u_{yy}v_{yy}) dx dy.$$

Let  $T_h = \{e\}$  be a uniform rectangular partition of  $\Omega$  and  $V_h \subset H_0^2(\Omega)$  be the Hermite bicubic finite element space associated with  $T_h$ . We use  $Q_p(e)$  represent the complete polynomial of degree  $p$ .  $v \in V_h$  if and only if

- 1)  $v|_e \in Q_3(e), \quad \forall e \in T_h;$
- 2)  $v, v_x, v_y, v_{xy}$  are continuous at the vertices and vanish at the vertices along  $\partial\Omega$ .

The finite element scheme is: find a pair of  $(\lambda_h, u_h)$ ,  $\lambda_h \in R$ ,  $u_h \in V_h$ ,  $\|u_h\|_0 = 1$ , such that

$$a(u_h, v) = \lambda_h(u_h, v), \quad \forall v \in V_h. \quad (2)$$

Let  $(\lambda, u)$  and  $(\lambda_h, u_h)$  denote the exact eigenpair of biharmonic operator and the associated approximate eigenpair in the finite element space  $V_h$ , respectively.  $R_h u \in V_h$  represents the Ritz-Galerkin projection of eigenfunction  $u$ .  $C > 0$  acts as a positive constant but it can take different value in this paper.  $C_\epsilon > 0$  is a positive number which depends on  $\epsilon$ . It is well known that<sup>[1,3,5]</sup>:

$$\lambda = \inf_{\phi \neq \theta, \phi \in H_0^2} \frac{\|\phi\|_a^2}{(\phi, \phi)}, \quad (3)$$

$$\lambda_h = \inf_{v \neq \theta, v \in V_h} \frac{\|v\|_a^2}{(v, v)}, \quad (4)$$

$$\|R_h u - u\|_2 \leq Ch^2 \|u\|_4, \quad (5)$$

where

$$\|w\|_a^2 = a(w, w), \quad \forall w \in H_0^2(\Omega).$$

### Proposition 1.

$$\begin{aligned} 0 \leq \lambda_h - \lambda &\leq Ch^4, \\ \|u_h - u\|_0 &\leq C_\epsilon h^{4-\epsilon}. \end{aligned}$$

where  $\epsilon > 0$  is an arbitrary small positive number.

*Proof.* First of all, we prove the first conclusion. Suppose  $(\lambda, u)$ , ( $\|u\|_0 = 1$ ) be an eigenpair of operator  $\Delta^2$ . Then we have

$$a(u, v) = \lambda(u, v), \quad \forall v \in H_0^2(\Omega). \quad (6)$$

obviously

$$(u, v) = 0 \iff a(u, v) = 0.$$

$\forall w \in H_0^2(\Omega)$ , let  $v = w - (w, u)u$ ,  $u_1 = (w, u)u$  then  $(v, u_1) = 0$  and  $a(v, u_1) = 0$ , namely

$$\|w\|_a^2 = \|u_1\|_a^2 + \|v\|_a^2.$$

Because  $\|u_1\|_a^2 = (w, u)^2 a(u, u) = \lambda(w, u)^2$ , we have the identity as below

$$\|w\|_a^2 = \lambda(u, w)^2 + \|w - (w, u)u\|_a^2, \quad (7)$$

therefore,  $\forall w \in H_0^2(\Omega)$ ,  $w \neq \theta$

$$\lambda \leq \frac{\|w\|_a^2}{\|w\|_0^2} \leq \lambda + \frac{\|w - (w, u)u\|_a^2}{\|w\|_0^2}. \quad (8)$$

Due to  $w - (w, u)u$  and space  $H_\lambda \equiv \{\alpha u : \alpha \in R\}$  are orthogonal for energy inner product  $a(\cdot, \cdot)$ , so

$$\|w - (w, u)u\|_a^2 \leq \|w - u\|_a^2$$

and

$$0 \leq \frac{a(w, w)}{(w, w)} - \lambda \leq \frac{\|w - u\|_a^2}{(w, w)}, \quad \forall w \in H_0^2(\Omega). \quad (9)$$

Note that  $V_h \subset H_0^2(\Omega)$ ,  $R_h u \subset V_h$ , we have

$$0 < \lambda \leq \frac{a(u_h, u_h)}{(u_h, u_h)} = \lambda_h \leq \frac{a(v, v)}{(v, v)}, \quad \forall v \in V_h \quad (10)$$

and

$$0 \leq \lambda_h - \lambda \leq \frac{a(R_h u, R_h u)}{(R_h u, R_h u)} - \lambda \leq \frac{\|R_h u - u\|_a^2}{(R_h u, R_h u)}. \quad (11)$$

From (11) and (5), for sufficient small  $h > 0$ , it is easy to gain our conclusion:

$$0 \leq \lambda_h - \lambda \leq Ch^4. \quad (12)$$

Secondly, for seeking the estimation of  $\|u_h - u\|_0$ , we consider the auxiliary problem: find  $w \in H_0^2(\Omega)$  such that

$$a(w, v) = (u_h, v), \quad \forall v \in H_0^2(\Omega). \quad (13)$$

Let  $\hat{u}$  and  $R_h \hat{u} \in V_h$  denote the exact solution and finite element solution, respectively. we can go the same way as the proof of Theorem 1<sup>[4]</sup> and get

$$\|u_h - u\|_0 \leq C\|\hat{u} - R_h \hat{u}\|_0. \quad (14)$$

Now introduce another auxiliary problem:  $\forall g \in H^{-1}(\Omega)$ , let  $r$  be the solution of problem:

$$\begin{cases} \Delta^2 r = g & \text{in } \Omega \\ r|_{\partial\Omega} = \frac{\partial r}{\partial n}|_{\partial\Omega} = 0 \end{cases}$$

then

$$\|r\|_3 \leq C\|g\|_{-1},$$

and

$$\|r\|_{4-\epsilon} \leq C_\epsilon \|g\|_0.$$

Using Nitsche technique, we have

$$\begin{aligned} |(\hat{u} - R_h \hat{u}, g)| &= |a(\hat{u} - R_h \hat{u}, r)| \\ &= |a(\hat{u} - R_h \hat{u}, r - r^I)| \\ &\leq C\|\hat{u} - R_h \hat{u}\|_2\|r - r^I\|_2 \\ &\leq C_\epsilon h^{4-\epsilon}\|\hat{u}\|_4\|r\|_{4-\epsilon} \\ &\leq C_\epsilon h^{4-\epsilon}\|\hat{u}\|_4\|g\|_0 \end{aligned}$$

where  $r^I \in V_h$  is the Hermite interpolation of  $r$ .

So, we get

$$\|\hat{u} - R_h \hat{u}\|_0 \leq C_\epsilon h^{4-\epsilon}. \quad (15)$$

Combine (14) and (15), we obtain

$$\|u_h - u\|_0 \leq C_\epsilon h^{4-\epsilon}. \quad (16)$$

**Proposition 2.** Suppose  $\epsilon > 0$  is an arbitrary small positive number, then

$$\|u_h - R_h u\|_2 \leq C_\epsilon h^{4-\epsilon}. \quad (17)$$

*Proof.*  $\forall v \in V_h$

$$\begin{aligned} |a(u_h - R_h u, v)| &= |a(u_h - u, v)| \\ &= |\lambda_h(u_h, v) - \lambda(u, v)| \\ &= |(\lambda_h - \lambda)(u_h, v) + \lambda(u_h - u, v)| \\ &\leq C(\lambda_h - \lambda + \|u_h - u\|_{-2})\|v\|_2. \end{aligned} \quad (18)$$

Take  $v = u_h - R_h u$  and we get our conclusion by (12), (16) and

$$\|u_h - u\|_{-2} \leq C\|u_h - u\|_0.$$

Finally, we present superconvergence approximations to the eigenvalue and eigenfunction.

Let  $T_{2h}$  is a uniform rectangular partition of  $\Omega$  with mesh parameter  $2h$ .  $T_h$  be obtained from dividing each element in  $T_{2h}$  into four congruent subelements, which belong to  $T_h$ . Define interpolation operator  $I^{2h}$  on  $T_{2h}$  satisfying

$$\begin{aligned} I^{2h}u|_e &\in Q_5(e) \quad \forall e \in T_{2h}, \\ I^{2h}u(A_i) &= u(A_i) \quad (I^{2h}u)_x(A_i) = u_x(A_i), \\ (I^{2h}u)_y(A_i) &= u_y(A_i) \quad (I^{2h}u)_{xy}(A_i) = u_{xy}(A_i). \quad i = 1, 2, \dots, 9 \end{aligned}$$

where  $A_i$  be the vertices of rectangular element in  $T_h$ .

**Lemma 1.** [2] Let  $T_h$  be a uniform mesh, then there holds the global superconvergence:

$$\|u - I^{2h}R_h u\|_2 \leq Ch^4 \|u\|_6$$

Now we can get our main result:

**Theorem 1.** If  $u \in H_0^2(\Omega) \cap H^6(\Omega)$ , then:

$$\|I^{2h}u_h - u\|_2 \leq C_\epsilon h^{4-\epsilon}, \quad (19)$$

$$0 \leq \bar{\lambda}_h - \lambda \leq C_\epsilon h^{8-\epsilon}, \quad (20)$$

where

$$\bar{\lambda}_h = \frac{a(I^{2h}u_h, I^{2h}u_h)}{(I^{2h}u_h, I^{2h}u_h)}.$$

*Proof.* Because of the identity:

$$I^{2h}u_h - u = I^{2h}(u_h - R_h u) + I^{2h}R_h u - u$$

so, from Proposition2 and Lemma1, we have

$$\begin{aligned} \|I^{2h}u_h - u\|_2 &\leq \|I^{2h}(u_h - R_h u)\|_2 + \|I^{2h}R_h u - u\|_2 \\ &\leq C\|u_h - R_h u\|_2 + \|I^{2h}R_h u - u\|_2 \\ &\leq C_\epsilon h^{4-\epsilon} \end{aligned}$$

Let

$$\bar{\lambda}_h = \frac{a(I^{2h}u_h, I^{2h}u_h)}{(I^{2h}u_h, I^{2h}u_h)}.$$

From (9), we have

$$0 \leq \bar{\lambda}_h - \lambda \leq \frac{\|I^{2h}u_h - u\|_a^2}{(I^{2h}u_h, I^{2h}u_h)}.$$

and through the continuity of  $a(\cdot, \cdot)$ , it is easy to obtain our conclusion:

$$0 \leq \bar{\lambda}_h - \lambda \leq C_\epsilon h^{8-\epsilon}.$$

**Acknowledgement.** The author thanks to Prof. Qun Lin for his guidance and Profs. Ningning Yan and Aihui Zhou for their discussions and suggestions.

## References

- [1] Qun Lin, Ningning Yan, High Efficiency FEM Construction and Analysis, Hebei Univ. Press, 1996.
- [2] Qun Lin, Ping Luo, Global superconvergence of Hermite bicubic element for the biharmonic equation, *Beijing Math.*, **1**:2 (1995), 52–64.
- [3] P.G. Ciarlet, J.L. Lions, Handbook of Numerical Analysis, Vol2, North-Holland, 1991.
- [4] Yang Yidu, Computable error bounds for an eigenvalue problem in the finite element method, *Chinese J. Num. Math. and Appl.*, **17**:1 (1995), 68–77.
- [5] Qun Lin, Qiding Zhu, The Preprocessing and Postprocessing for FEM, Shanghai Sci. and Tec. Publishers, 1994.
- [6] Yang Yidu, A posteriori error estimates in Adini finite element for eigenvalue problems, Preprint.
- [7] Qun Lin, Ningning Yan, Aihui Zhou, A rectangle test for interpolated finite elements, Pro. of Sys. Sci. and Sys. Engng, Great wall Culture publish Co., Hong Kong, 1991, 217–229.
- [8] R. Rannacher, Nonconforming finite element methods for eigenvalue problems in linear plate theory, *Numer. Math.*, **33** (1979), 23–42.