

# Oscillation Theory of $h$ -fractional Difference Equations\*

Fanfan Li<sup>1</sup> and Zhenlai Han<sup>1,†</sup>

**Abstract** In this paper, we initiate the oscillation theory for  $h$ -fractional difference equations of the form

$$\begin{cases} {}_a\Delta_h^\alpha x(t) + r(t)x(t) = e(t) + f(t, x(t)), & t \in \mathbb{T}_h^a, \quad 1 < \alpha < 2, \\ x(a) = c_0, \quad \Delta_h x(a) = c_1, \quad c_0, c_1 \in \mathbb{R}, \end{cases}$$

where  ${}_a\Delta_h^\alpha$  is the Riemann-Liouville  $h$ -fractional difference of order  $\alpha$ ,  $\mathbb{T}_h^a := \{a + kh, k \in \mathbb{Z}^+ \cup \{0\}\}$ , and  $a \geq 0$ ,  $h > 0$ . We study the oscillation of  $h$ -fractional difference equations with Riemann-Liouville derivative, and obtain some sufficient conditions for oscillation of every solution. Finally, we give an example to illustrate our main results.

**Keywords**  $h$ -difference equations, Oscillation, Fractional.

**MSC(2010)** 34A10, 34C10, 26A33.

## 1. Introduction

The fractional calculus (calculus with derivatives of arbitrary order) is an important research field in several different areas such as physics (including classical and quantum mechanic as well as thermodynamics), chemistry, biology, economic, and control theory [7, 10, 11, 13–15]. Fractional difference equations, which is the discrete counterpart of the corresponding fractional differential equations, have recently gained an intensive interest among researchers in the last years. The discrete fractional calculus has been developed much after the appearance of time scale calculus [6]. The researchers started to use delta and nabla analysis by employing the jumping operators  $\sigma$  and  $\rho$ .

Computer simulations show that the time scale  $(h\mathbb{Z})_a$  is particularly interesting because when  $h$  tends to zero one recovers previous fractional continuous-time results [5, 8]. However, the development of the qualitative features of these type of equations is still considered to be at its first stage of progress. Therefore, it is pertinent to develop the oscillation theory of fractional  $h$ -difference equations. The last few years, there are new papers which have studied oscillation of fractional difference equations, see the papers [1, 2, 4, 12].

---

<sup>†</sup>the corresponding author.

Email address: lifanfan1994@163.com(F. Li), hanzhenlai@163.com(Z. Han)

<sup>1</sup>School of Mathematical Sciences, University of Jinan, Jinan, Shandong 250022, China

\*The authors were supported by Shandong Provincial Natural Science Foundation (ZR2017MA043).

Marian et al. [12] studied the oscillation criteria for forced nonlinear fractional difference equations of the form

$$\begin{cases} \Delta^\alpha x(t) + f_1(t, x(t+a)) = v(t) + f_2(t, x(t+a)), & t \in \mathbb{N}_0, \quad 0 < \alpha \leq 1, \\ \Delta^{\alpha-1} x(t)|_{t=0} = x_0, \end{cases}$$

where  $\Delta^\alpha$  denotes the Riemann-Liouville like discrete fractional difference operator of order  $\alpha$ .

Abdalla et al. [1] studied the oscillation of solutions of nonlinear forced fractional difference equations of the form

$$\begin{cases} \nabla_{a(q)-1}^q x(t) + f_1(t, x(t)) = r(t) + f_2(t, x(t)), & t \in \mathbb{N}_{a(q)}, \\ \nabla_{a(q)-1}^{-(m-q)} x(t)|_{t=a(q)} = x(a(q)) = c, & c \in \mathbb{R}, \end{cases}$$

where  $q > 0$ ,  $m = [q] + 1$ ,  $m \in \mathbb{N}$ ,  $[q]$  is the greatest integer less than or equal to  $q$ ,  $\nabla_{a(q)-1}^{-q}$  and  $\nabla_{a(q)-1}^q$  are the Riemann-Liouville sum and difference operators, respectively.

Following this trend and there are no results available in the literature regarding the oscillation of solutions of  $h$ -fractional difference equations, we are concerned equations of the form

$$\begin{cases} {}_a\Delta_h^\alpha x(t) + r(t)x(t) = e(t) + f(t, x(t)), & t \in \mathbb{T}_h^a, \quad 1 < \alpha < 2, \\ x(a) = c_0, \quad \Delta_h x(a) = c_1, & c_0, c_1 \in \mathbb{R}, \end{cases} \quad (1.1)$$

where  ${}_a\Delta_h^\alpha$  is the Riemann-Liouville  $h$ -fractional difference of order  $\alpha$ ,  $\Delta_h$  is the forward  $h$ -difference operator,  $\mathbb{T}_h^a := \{a + kh, k \in \mathbb{Z}^+ \cup \{0\}\}$ ,  $a \geq 0$ ,  $h > 0$  and  $e(t)$  is a continuous function. The problems will be studied under the following assumptions:

- (H1)  $r(t)$  is a positive real-valued continuous function on  $\mathbb{R}$ ;
- (H2)  $x(t)f(t, x(t)) > 0$ ,  $x(t) \neq 0$ ,  $t \in \mathbb{T}_h^a$ ;
- (H3)  $|f(t, x(t))| \leq p(t)|x(t)|^\gamma$ ,  $x(t) \neq 0$ ,  $t \geq a$ ;
- (H4)  $|f(t, x(t))| \geq p(t)|x(t)|^\gamma$ ,  $x(t) \neq 0$ ,  $t \geq a$ ,

where  $p(t) \in C(\mathbb{T}_h^a, \mathbb{R}^+)$  and  $\gamma$  is a the quotient of two positive odd numbers.

We only consider these solutions of (1.1) which exist on  $\mathbb{T}_h^a$ . If  $x(t)$  satisfies (1.1) on  $\mathbb{T}_h^a$ , then the function  $x(t)$  is called a solution of (1.1). A solution  $x(t)$  of (1.1) is said to be oscillatory if it is neither eventually positive nor eventually negative, otherwise it is called nonoscillatory. The equation itself is called oscillatory if all of its solutions are oscillatory.

The paper is organized as follows. In Section 2, we present some basic definitions and useful results from the theory of discrete fractional calculus which we rely in the later section. In Section 3, we intend to use the inequalities to obtain some sufficient conditions for oscillation for oscillation of every solution of (1.1). In Section 4, we give an example to illustrate our results.

## 2. Preliminaries

Before stating and proving our results, we introduce some definitions and notations. Let  $\mathbb{T}_h^a := \{a + kh, k \in \mathbb{Z}^+ \cup \{0\}\}$ , where  $a \geq 0$ ,  $h > 0$ . Let us denote by  $\mathcal{F}_{\mathbb{T}}$  the set of real valued functions defined on  $\mathbb{T}_h^a$ ,  $\sigma(t) = t + h$  and  $\rho(t) = t - h$ .