

# COERCIVITY OF THE SINGLE LAYER HEAT POTENTIAL

DOUGLAS N. ARNOLD    PATRICK J. NOON

*(Department of Mathematics, University of Maryland, College park, MD20742, USA)*

## Abstract

The single layer heat potential operator,  $\mathcal{K}$ , arises in the solution of initial-boundary value problems for the heat equation using boundary integral methods. In this note we show that  $\mathcal{K}$  maps a certain anisotropic Sobolev space isomorphically onto its dual, and, moreover, satisfies the coercivity inequality  $\langle \mathcal{K}_{q,q} \rangle \geq c \|q\|^2$ . We thereby establish the well-posedness of the operator equation  $\mathcal{K}_q = f$  and provide a basis for the analysis of the discretizations.

## §1. Introduction

If  $u(x, t)$  solves the homogeneous heat equation for  $x$  in a smoothly bounded domain  $\Omega$  in  $\mathbb{R}^3$  and  $t > 0$  and vanishes when  $t = 0$ , then  $u$  may be expressed in terms of its Cauchy data on  $\Gamma \times \mathbb{R}_+$  as

$$u(x, t) = \int_0^t \int_{\Gamma} \left[ \frac{\partial u}{\partial n}(y, s) K(x - y, t - s) - \frac{\partial K}{\partial n}(x - y, t - s) u(y, s) \right] dy ds$$

where  $\Gamma = \partial\Omega$  and  $K(x, t)$  denotes the fundamental solution to the heat equation,

$$K(x, t) = \begin{cases} \frac{\exp(-|x|^2/4t)}{(4\pi t)^{3/2}}, & x \in \mathbb{R}^3, t > 0, \\ 0, & x \in \mathbb{R}^3, t \leq 0. \end{cases}$$

Letting  $x$  tend to  $\Gamma$  and employing a jump relation [5, p. 137] leads to a mixed Volterra-Fredholm integral equation relating the values of  $u$  and  $\partial u/\partial n$  on the boundary of the space-time cylinder. Together with given boundary conditions, this integral equation can be used to solve for the Cauchy data of  $u$  and then  $u$  can be determined globally from the representation formula. Thus the solution of



boundary value problems for the heat equation can be reduced to the solution of an integral equation posed on the boundary of the cylinder. This approach is employed with increasing frequency for the numerical solution of transient heat conduction problems [ 2, 4, 7, 9, 12].

The form of the integral equation to be solved depends on the boundary value problem considered. If Neumann or Robin conditions are specified for  $u$ , then the unknown Dirichlet data of  $u$  is determined by an integral equation of the second kind. This second kind equation has been studied in depth by Pogorzelski [ 10, 11] and his results have recently been applied to the analysis of numerical methods based on this boundary integral formulation [3]. On the other hand, if  $u$  solves a Dirichlet problem, then the representation formula leads to an integral equation of the first kind of the form

$$K_q(x, t) := \int_0^t \int_{\Gamma} q(y, s)K(x - y, t - s)dyds = f(x, t), \quad x \in \Gamma, t > 0, \quad (1)$$

where  $q$  is the unknown flux and  $f$  is known. Our aim in this note to prove a simple coercivity estimate for this operator which implies the well-posedness of the integral equation and provides a basis for the analysis of discretizations. It is appropriate to work with the anisotropic Sobolev space  $H^{r,s}(S, I) := L^2(I, H^r(S)) \cap H^s(I, L^2(S))$  and its dual  $H^{-r,-s}(S, I)$  ( $r, s \geq 0, S$  a smooth submanifold of  $\mathbb{R}^3, I$  an interval.) Cf. [6, Ch. 4]. Our major result may be stated as follows:

**Theorem 1.** *The single layer heat potential operator  $K$  defines an isomorphism from the Hilbert space  $H^{-1/2,-1/4}(\Gamma, \mathbb{R}_+)$  onto its dual. Moreover the coercivity estimate*

$$\langle K_{q,q} \rangle \geq c \|q\|_{-1/2,-1/4}^2 \quad \text{for all } q \in H^{-1/2,-1/4}(\Gamma, \mathbb{R}_+)$$

holds for some positive constant  $c$ .

*Remarks.* 1. Nedelec and Planchard [8] proved a similar result for the electrostatic single layer potential equation for the Laplace equation. It is remarkable that such a coercivity estimate, typical of elliptic operators, also holds for the single layer heat potential. 2. R. Brown [1] has recently shown that  $K$  maps  $L^2(\Gamma \times \mathbb{R}_+)$  isomorphically onto  $H^{1,1/2}(\Gamma, \mathbb{R}_+)$ . In the present context, this may be viewed as one of many possible regularity results.

### §2. Factorization of the heat potential

For  $u$  in the Schwartz class  $S(\mathbb{R}^3 \times \mathbb{R})$  set

$$\begin{aligned} \|u\|_{W^{1,0}}^2 &= \int_{\mathbb{R}} \int_{\mathbb{R}^3} |\xi|^2 |\hat{u}(\xi, \tau)|^2 d\xi d\tau, \\ \|u\|_{W^{1,1/2}}^2 &= \int_{\mathbb{R}} \int_{\mathbb{R}^3} (|\xi|^2 + |\tau|) |\hat{u}(\xi, \tau)|^2 d\xi d\tau, \\ \|u\|_{\mathcal{V}}^2 &= \int_{\mathbb{R}} \int_{\mathbb{R}^3} (|\xi|^2 + |\tau|^2 |\xi|^{-2}) |\hat{u}(\xi, \tau)|^2 d\xi d\tau, \end{aligned}$$